PLANT FACT SHEET

Gelidium latifolium (Greville)Bornet & Thuret

GENUS *Gelidium* **FAMILY** Gelidiaceae

COMMON NAMES Jelly plant

EDIBLE PARTS AND USES

Edible parts: Seaweed, Frond, Algae

It is dried, bleached and boiled to improve the flavour.
It is a main source of agar.

DESCRIPTION

A seaweed.

HABITAT

It grows in temperate and tropical places.

REFERENCES

Brouk, B., 1975, Plants Consumed by Man. Academic Press, London. p 253

Burkill, I.H., 1966, A Dictionary of the Economic Products of the Malay Peninsula. Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia. Vol 1 (A-H) p 1081

http://www.seavegetables.com

Hu, Shiu-ying, 2005, Food Plants of China. The Chinese University Press. p 261

Kiple, K.F. & Ornelas, K.C., (eds), 2000, The Cambridge World History of Food. CUP p 237

Subba Rao, G.N., 1965, Uses of seaweed directly as human food. Indo-Pacific Fisheries Council Regional Studies 2:1-32

Surey-Gent, S. & Morris G., 1987, Seaweed. A User's Guide. Whirret Books. London. p 62, 135

Zaneveld, J.S., 1955, Economic marine algae of tropical South and East Asia and their utilization. Indo-Pacific Special Publications, No 3 Bangkok.

Zemke-White, W. L. & Ohno, M., 1999, World seaweed utilisation: An end-of-century summary. Journal of Applied Phycology 11: 369-376

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/45083dd4-b34f-4616-8b52-42aeac498428