



PLANT FACT SHEET

Vigna subterranea (L.) Verdc.

GENUS *Vigna* FAMILY Fabaceae

COMMON NAMES Bambara groundnut

EDIBLE PARTS AND USES

Edible parts: Seeds, Leaves, Pods, Vegetable, Nuts, Seeds - coffee

Seeds can be eaten fresh or roasted while immature. Mature seeds are hard so must be boiled before being used in cooking. Pre-soaking seeds and cooking with potash makes cooking easier.

Young pods are cooked and used as a vegetable. They are used in stews.

Seeds can be dried and made into flour and used for baking. They can be popped like corn.

The seeds are roasted as a coffee substitute.

The leaves can be eaten.

DESCRIPTION

An annual plant. It can either form a bunchy bush or be a trailing plant. Often the creeping stems are near ground level. It often appears as if bunched leaves arise from branched stems near ground level. It has a well developed taproot. The leaves have 3 leaflets. The leaf stalk is erect and thickened near the base. The end leaflet is slightly larger than the side leaflets. Leaflets are about 6 cm long by 3 cm across. The flowers are yellowish-white. They occur in pairs. The fruit are pods which are round and with one seed. Some kinds have 3 seeds. This pod develops under the ground on a long stalk. The flower/fruit stalk elongates after being fertilised and pushes into the soil. The seeds are hard and are of many colours. Pods can be 3.7 cm long.

HABITAT

It is a tropical plant. It can grow in hot climates. It can also grown on poor soils. It does best with moderate rainfall and sunshine. It can tolerate drought. Long day-lengths can reduce or prevent pod development in some kinds. It suits semi-arid regions. In Kenya it grows from sea level to 1,550 m altitude. It can grow in arid places.

PRODUCTION

Cultivation

Plants are grown from seed. Plants are often put in rows 50 cm apart and with 15 cm spacing between plants. Ridges are formed to enable the pods to penetrate the soil. It is mostly grown intercropped with other plants. Soil should be light and friable and the seed bed loose and fine. Normally the whole plant is pulled up for harvesting. Any pods which become detached are harvested by hand. Pods are dried in the air before threshing.

Harvesting and Seed Production

Pods reach their maximum size 30 days after being fertilised and seeds develop over the next 10 days. Pods mature in 4 months from planting. Yields of 500-1,000 kg of dried nuts per hectare are produced. Higher yields are possible.

NOTES

On average, the seeds were found to contain 63% carbohydrate, 19% protein and 6.5% oil.

PICTURES



Credit: Food Plants International (Bruce French)



Credit: Food Plants International (Bruce French)

REFERENCES

- Abbiw, D.K., 1990, Useful Plants of Ghana. West African uses of wild and cultivated plants. Intermediate Technology Publications and the Royal Botanic Gardens, Kew. p 31
- Akinola, R., et al, 2020, A Review of Indigenous Food Crops in Africa and the Implications for more Sustainable and Healthy Food Systems. Sustainability 2020, 12, 3493;
- Ambasta, S.P. (Ed.), 2000, The Useful Plants of India. CSIR India. p 685 (As Voandzeia subterranea)
- Asfaw, Z. and Tadesse, M., 2001, Prospects for Sustainable Use and Development of Wild Food Plants in Ethiopia. Economic Botany, Vol. 55, No. 1, pp. 47-62
- Baa-Poku, F. & Asante, I. K., 2020, Use of Neglected and Underutilized Crop Species (NUCS) in some farming communities in the Upper Afram and Upper Dayi River Basins in Ghana. International Journal of Technology and Management Research 5 (2): 48-63
- Baidu-Forson, J.J., et al ,2014,. Assessment of agrobiodiversity resources in the Borotse flood plain, Zambia. CGIAR Research Program on Aquatic Agricultural Systems. Penang, Malaysia. Working Paper: AAS-2014-12.

- Brink, M., Ramolemana, G.M. & Sibuga, K.P., 2006. *Vigna subterranea* (L.) Verdc. [Internet] Record from Protabase. Brink, M. & Belay, G. (Editors). PROTA (Plant Resources of Tropical Africa), Wageningen, Netherlands. < <http://database.prota.org/search.htm>>. Accessed 23 October 2009
- Burkill, H. M., 1985, The useful plants of west tropical Africa, Vol. 3. Kew.
- Burkill, I.H., 1966, A Dictionary of the Economic Products of the Malay Peninsula. Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia. Vol 2 (I-Z) p 2291 (As *Voandzeia subterranea*)
- Cobley, L.S. (rev. Steele, W.M.) 2nd Ed., 1976, An Introduction to the Botany of Tropical Crops. Longmans. p 101 (As *Voandzeia subterranea*)
- Dakora, F. D., 2013, Biogeographic Distribution, Nodulation and Nutritional Attributes of Underutilized Indigenous Agrican Legumes. Acta Horticulturae Number 979 Vol. 1. p 53
- Dalziel, J. M., 1937, The Useful plants of west tropical Africa. Crown Agents for the Colonies London.
- Dansi, A., et al, 2012, Diversity of the Neglected and Underutilized Crop Species of Importance in Benin. The Scientific World Journal. Volume 2012, Article ID 932947, 19 pages
- Facciola, S., 1998, Cornucopia 2: a Source Book of Edible Plants. Kampong Publications, p 114 (As *Voandzeia subterranea*)
- FAO, 1988, Traditional Food Plants, FAO Food and Nutrition Paper 42. FAO Rome p 507
- FAO, 2024, Compendium of forgotten foods in Africa - A companion publication for Integrating Africa's forgotten foods for better nutrition. Accra. <https://doi.org/10.4060/cc5044en> p 46
- Fowler, D. G., 2007, *Zambian Plants: Their Vernacular Names and Uses*. Kew. p 41
- Fox, F. W. & Young, M. E. N., 1982, *Food from the Veld*. Delta Books. p 223 (As *Voandzeia subterranea*)
- Grivetti, L. E., 1980, Agricultural development: present and potential role of edible wild plants. Part 2: Sub-Saharan Africa, Report to the Department of State Agency for International Development. p 32
- Hedrick, U.P., 1919, (Ed.), *Sturtevant's edible plants of the world*. p 688 (As *Voandzeia subterranea*)
- Heller, J., F. Begemann and J. Mushonga, editors. 1997. Bambara groundnut (*Vigna subterranea* (L.) Verdc.). Promoting the conservation and use of underutilized and neglected crops. 9. Proceedings of the workshop on Conservation and Improvement of Bambara Groundnut (*Vigna subterranea* (L.) Verdc.), 14-16 November 1995, Harare, Zimbabwe. Institute of Plant Genetics and Crop Plant Research, Gatersleben/Department of Research & Specialist Services, Harare/International Plant Genetic Resources Institute, Rome, Italy.
- ILDIS Legumes of the World <http://www.ildis.org/Legume/Web>
- Jacquat, C., 1990, *Plants from the Markets of Thailand*. D.K. Book House p (As *Voandzeia subterranea*)
- Jardin, C., 1970, List of Foods Used In Africa, FAO Nutrition Information Document Series No 2.p 110 (As *Voandzeia subterranea*)
- Kays, S. J., and Dias, J. C. S., 1995, Common Names of Commercially Cultivated Vegetables of the World in 15 languages. Economic Botany, Vol. 49, No. 2, pp. 115-152
- Kermath, B. M., et al, 2014, *Food Plants in the Americas: A survey of the domesticated, cultivated and wild plants used for Human food in North, Central and South America and the Caribbean*. On line draft. p 911
- Kew Bull. 35:474. 1981
- Kiple, K.F. & Ornelas, K.C., (eds), 2000, *The Cambridge World History of Food*. CUP p 1726 (As *Voandzeia subterranea*)
- Koukou, E., et al, 2022, Local knowledge of agricultural biodiversity and food uses in two agroecological zones of southern Benin. *Ethnobotany Research and Applications* 23:29 (2022) p 10
- Latham, P., 2004, *Useful Plants of Bas-Congo province*. Salvation Army & DFID p 289
- Long, C., 2005, *Swaziland's Flora - siSwati names and Uses* <http://www.sntc.org.sz/flora/> (As *Voandzeia subterranea*)
- Macmillan, H.F. (Revised Barlow, H.S., et al), 1991, *Tropical Planting and Gardening*. Sixth edition. Malayan Nature Society. Kuala Lumpur. p 350

Full reference list (38 more) at <https://www.echocommunity.org/en/resources/4a3764a0-b7a2-45b4-8ad0-6fa85d509464>

PROVIDED BY Food Plants International

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET <https://www.echocommunity.org/en/resources/4a3764a0-b7a2-45b4-8ad0-6fa85d509464>