PLANT FACT SHEET

Caulerpa sertularioides (Gmelin) Howe

GENUS *Caulerpa* **FAMILY** Caulerpaceae

COMMON NAMES Feather Caulerpa

EDIBLE PARTS AND USES

Edible parts: Algae, Seaweed

It is eaten raw or as a salad. It is eaten with lemon juice and coconut milk.

DESCRIPTION

A green seaweed. The branches are erect. They are feather like. They are fine and not flattened. They are usually in 2 rows but can be in 3 rows.

HABITAT

It is a tropical plant. It grows just below the tidal zones. It grows on sheltered and partly exposed shores.

REFERENCES

Ambasta S.P. (Ed.), 2000, The Useful Plants of India. CSIR India. p 112

Burkill, H. M., 1985, The useful plants of west tropical Africa, Vol. 5. Kew.

Cribb, A.B., 1996, Seaweeds of Queensland A Naturalist's Guide. The Queensland Naturalists' Club Handbook No. 2. p 21

Daillande, C. de, et al, 2016, Caulerpa consumption, nutritional value and farming in the Indo-Pacific region. Journal of Applied Phycology. July 2016

http://www.seavegetables.com

Novaczek, I., 2001, A Guide to the Common Edible and Medicinal Plants of the Pacific Islands. SPC. p 5

Ostraff, M., 2003, Contemporary Uses of Limu (marine algae) in Tonga. Ph. D. thesis. p 81

Wang, Wei-Lung and Chiang, Young-Meng, 1994, Potential Economic Seaweeds of Hengchun Peninsula, Taiwan, Economic Botany, Vol. 48, No. 2, pp. 182-189

Womersley, H.B. S. & Bailey, A., 1970, Marine algae of the Solomon Islands. Philosophical Transactions of the Royal Society of London. B. Biological Sciences 259:257-352

www.algaebase/org

Zemke-White, W. L. & Ohno, M., 1999, World seaweed utilisation: An end-of-century summary. Journal of Applied Phycology 11: 369-376

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/4f4fb8fd-52f2-4600-84e9-4154d5faafc8