PLANT FACT SHEET

# *Chaetomorpha antennina* (Bory de Saint-Vincent) Kuetzing

**GENUS** *Chaetomorpha* **FAMILY** Cladophoraceae

## EDIBLE PARTS AND USES

**Edible parts:** Fronds, Seaweed

## DESCRIPTION

A seaweed. The fronds are not branched. They are stiff and bright green. They are attached in clumps. The large cells making up the frond filaments can be seen with the eye.

## HABITAT

A tropical plant. It grows in the middle of the tidal zone. It can be on steep rocks on exposed shores.

## REFERENCES


Burkill, I.H., 1966, A Dictionary of the Economic Products of the Malay Peninsula. Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia. Vol 1 (A-H) p 525

Cribb, A.B., 1996, Seaweeds of Queensland A Naturalist's Guide. The Queensland Naturalists' Club Handbook No. 2. p 21

Kiple, K.F. & Ornelas, K.C., (eds), 2000, The Cambridge World History of Food. CUP p 234

Wang, Wei-Lung and Chiang, Young-Meng, 1994, Potential Economic Seaweeds of Hengchun Peninsula, Taiwan, Economic Botany, Vol. 48, No. 2, pp. 182-189

Womersley, H.B. S. & Bailey, A., 1970, Marine algae of the Solomon Islands. Philosophical Transactions of the Royal Society of London. B. Biological Sciences 259:257-352

Zaneveld, J.S., 1950, The economic marine algae of Malaysia and their applications. Proceedings of the Indo-Pacific Fisheries Council, 107-14



**PROVIDED BY** **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International



**LICENCE** CC BY-NC-SA 3.0


**ONLINE FACT SHEET** https://www.echocommunity.org/en/resources/72f3f954-a409-4838-91a3-40fd725a4089