PLANT FACT SHEET

Puschkinia scilloides Adams

GENUS *Puschkinia* **FAMILY** Asparagaceae

COMMON NAMES Striped squill

EDIBLE PARTS AND USES

Edible parts: Leaves, Flowers

It is cooked by boiling or used in soups.
The leaves are also eaten raw in salads.

DESCRIPTION

A herb. It grows from a bulb and re-grows each year. It grows in a small clump. It grows 20 cm tall. The flowers are bell shaped and pale blue with darker blue stripes.

HABITAT

It is a temperate plant.

PRODUCTION

Cultivation

It can be grown by seed. It can also be grown by bulbs.

NOTES

Also put in the family Hyacinthaceae.

REFERENCES

Cakir, E. A., 2017, Traditional knowledge of wild edible plants of Iğdır Province (East Anatolia, Turkey). Acta Soc Bot Pol. 2017;86(4):3568

Ertug, F, Yenen Bitkiler. Resimli Türkiye Florası -I- Flora of Turkey - Ethnobotany supplement (Also as Puschkinia scilloides var. libanotica)

Hovsepyan, R., et al, 2016, Food as a marker for economy and part of identity: traditional vegetal food of Yezidis and Kurds in Armenia. Journal of Ethnic Foods. 3:32-41

http://www.botanic-gardens-ljubljana.com/en/plants

Ibadullayeva, S., et al, 2011, Protection of some rare and endangered vegetable plants in the flora of the Nakhichevan AR. International Journal of Biodiversity and Conservation Vol. 3(6), pp. 224-229

Kayiran, S. D., & Ozkan, E. E., 2017, The ethnobotanical uses of Hyacinthaceae species growing in Turkey and a review of pharmacological activities. Indian Journal of Traditional Knowledge Vol. 16(2), April 2017, pp. 243-250

Nanagulyan, S., et al, 2020, Wild plants and fungi sold in the markets of Yerevan (Armenia). Journal of Ethnobiology and Ethnomedicine 16:26

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/74086dcc-5c99-44f2-8128-c9a478beeb7e