PLANT FACT SHEET

Beta vulgaris var. altissima L.; Doll

GENUS *Beta* **FAMILY** Amaranthaceae

COMMON NAMES Sugarbeet

EDIBLE PARTS AND USES

Edible parts: Sugar - roots, Spice, Flavour, Leaves

The leaves are used for sarma in Turkey. They are rolled around a filling of rice or minced meat.

DESCRIPTION

A herb.

HABITAT

It is a temperate plant.

NOTES

Probably all Beta are one species and 2 main forms - Cicla - for leaves, and Contiva - for roots.
Also put in the family Chenopodiaceae.

REFERENCES

Brouk, B., 1975, Plants Consumed by Man. Academic Press, London. p 247, 364

Campbell, G.K.G., 1979, Sugar beet, in Simmonds N.W.,(ed), Crop Plant Evolution. Longmans. London. p 25

Dogan, Y., et al, 2015, Of the importance of a leaf: the ethnobotany of sarma in Turkey and the Balkans. Journal of Ethnobiology and Ethnomedicine, 11:56

Jardin, C., 1970, List of Foods Used In Africa, FAO Nutrition Information Document Series No 2.p 21

Kiple, K.F. & Ornelas, K.C., (eds), 2000, The Cambridge World History of Food. CUP p 434

Plants for a Future database, The Field, Penpol, Lostwithiel, Cornwall, PL22 0NG, UK. http://www.scs.leeds.ac.uk/pfaf/

Rhein. Fl. 293. 1843

Vorstenbosch, T., et al, 2017, Famine food of vegetal origin consumed in the Netherlands during World War II. Journal of Ethnobiology and Ethnomedicine (2017) 13:63

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/7eae0103-2b77-4da5-acba-50fa969f2087