PLANT FACT SHEET

Amelanchier interior Nielson

GENUS *Amelanchier* **FAMILY** Rosaceae

COMMON NAMES Inland Service-berry, Pacific Serviceberry

EDIBLE PARTS AND USES

Edible parts: Fruit

The fruit is eaten raw or cooked.

DESCRIPTION

A tree to 9 m high.

HABITAT

It is a temperate plant.

PRODUCTION

Cultivation

They are grown from ripened seed. It can be grown from seed, layering or suckers. Seed can take 18 months to germinate and layers can take 18 months to form roots.

NOTES

There are about 25-30 Amelanchier species. Possibly a natural hybrid involving A. laevis, A. arborea, and A. spicata, or A. sanguinea

REFERENCES

Amer. Midl. Naturalist 22:185, t. 13. 1939

Fisk, J. R. & Hoover, E., 2015, Wild Fruits of Minnesota. A Filed Guide. University of Minnesota p 8

Grandtner, M. M., 2008, World Dictionary of Trees. Wood and Forest Science Department. Laval University, Quebec, Qc Canada. (Internet database http://www.WDT.QC.ca)

Kermath, B. M., et al, 2018, Food Plants in the Americas: A Survey of the Domesticated, Cultivated and Wild Plants Used for Human Food in North, Central and South America and the Caribbean. p 71

Plants for a Future database, The Field, Penpol, Lostwithiel, Cornwall, PL22 0NG, UK. http://www.scs.leeds.ac.uk/pfaf/.

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/84117488-22bb-4ae8-8b9b-6b1b0f4e4ee5