PLANT FACT SHEET

Mandevilla tenuifolia (J. C. Mikan) Woodson

GENUS *Mandevilla* **FAMILY** Apocynaceae

COMMON NAMES Cauhy

EDIBLE PARTS AND USES

Edible parts: Rhizome, Tubers, Root, Stem

The tuber is cooked and eaten. It is also eaten raw and preserved in pickles.

DESCRIPTION

A herb or vine. It grows 30 cm tall. The tuber is the size of a turnip. The stems are slender. The leaves are narrow and 3-12 cm long by 2-5 mm wide.

HABITAT

It is a tropical plant.

REFERENCES

Allorge-Boiteau, L., Flora of the Guianas. 140 Apocynaceae p 149

Bussmann, R. W., (Ed.), 2020, Biodiverse food plants in the semiarid region of Brazil have unknown potential: A systematic review. Plos ONE 15(5): e0230936

Cruz, M. P., et al, 2014, Local perceptions and consumption of native wild edible plants from seasonal dry forests in Brazil. Journal or Ethnobiology and Ethnomedicine 10:45

Fagg, C. W. et al, 2015, Useful Brazilian plants listed in the manuscripts and publications of the Scottish medic and naturalist George Gardner (1812–1849). Journal of Ethnopharmacology 161 (2015) 18–29

Gomes, L.C.A., et al, 2023, Patterns of use of wild food plants by Brazilian local communities: systematic review and meta-analysis. Journal of Ethnobiology and Ethnomedicine. 19:47 p 7

Hedrick, U.P., 1919, (Ed.), Sturtevant's edible plants of the world. p 279 (As Dipladenia tenuifolia)

Nascimento, V. T. D., et al, 2012, Famine Foods of Brazil's Seasonal Dry Forests: Ethnobotanical and Nutritional Aspects. Economic Botany.66 (1), pp 22-34

Nascimento, V. T. D., et al, 2013, Knowledge and Use of Wild Food Plants in Areas of Dry Seasonal Forests in Brazil. Ecology of Food and Nutrition, 52:317–343

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/84f0c3d1-0a1d-4da2-a875-1ae439826495