PLANT FACT SHEET

Urochloa stigmatisata (Mez) K.M.Ibrahim & P.M.Peterson.

GENUS *Urochloa* **FAMILY** Poaceae

EDIBLE PARTS AND USES

Edible parts: Seeds, Cereal

The seeds are cooked and eaten.
They are also used to make a drink.

DESCRIPTION

A small grass. It often lies along the ground. It grows each year from seeds.

HABITAT

A tropical plant. It grows in disturbed soils in drier regions of West Africa.

NOTES

There are about 100 Brachiaria species.

REFERENCES

Abbiw, D.K., 1990, Useful Plants of Ghana. West African uses of wild and cultivated plants. Intermediate Technology Publications and the Royal Botanic Gardens, Kew. p 25

Bircher, A. G. & Bircher, W. H., 2000, Encyclopedia of Fruit Trees and Edible Flowering Plants in Egypt and the Subtropics. AUC Press. p 63

Burkill, H. M., 1985, The useful plants of west tropical Africa, Vol. 3. Kew.

Busson, 1965,

Gallagher, D. E., 2010, Farming beyond the escarpment: Society, Environment, and Mobility in Precolonial Southeastern Burkina Faso. PhD University of Michigan.

Hahn, K., et al, 2018, The use of wild plants for food: a national scale analysis for Burkina Faso (West Africa). FVSS FoodPlants Supplement

Jardin, C., 1970, List of Foods Used In Africa, FAO Nutrition Information Document Series No 2.p 1

Peters, C. R., O'Brien, E. M., and Drummond, R.B., 1992, Edible Wild plants of Sub-saharan Africa. Kew. p 19

World Checklist of Useful Plant Species 2020. Royal Botanic Gardens, Kew

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/875f669d-819d-4368-b27e-3139437aa8a2