PLANT FACT SHEET

Diplazium stoliczkae Beddome

GENUS *Diplazium* **FAMILY** Athyriaceae

COMMON NAMES Kalo nyuro

EDIBLE PARTS AND USES

Edible parts: Leaves, Fronds

The young fronds are cooked as a vegetable.

DESCRIPTION

A fern. It has a smooth leaf stalk and rachis. The fronds are 10-90 cm long by 15-30 cm wide. There are many leaflets (pinnae) and they taper towards the tip. The cluster of spores are in a line.

HABITAT

It is a temperate plant. In Nepal it grows from 1500-4000 m altitude.

PRODUCTION

Cultivation

Plants can be grown from spores or by splitting the rhizome.

NOTES

There are about 300-600 Diplazium species. They have also been put in the families Aspleniaceae and Woodsiaceae.

REFERENCES

Joshi, N., et al, 2007, Traditional neglected vegetables of Nepal: Their sustainable utilization for meeting human needs. Tropentag 2007. Conference on International Agricultural Research for Development.

Joshi, K. and Joshi, A. R., 2008, Ethnobotanical Studies on Some Lower Plants of the Central Development Region, Nepal. Ethnobotanical Leaflets 12:832-40

Manandhar, N.P., 2002, Plants and People of Nepal. Timber Press. Portland, Oregon. p 204

Ojha, R. & Devkota, H. P., 2021, Edible and Medicinal Pteridophytes of Nepal: A Review. Ethnobotany Research & Applications 22:16

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/8c7c57c5-de4e-4002-a6ee-9800c0cabc7f