PLANT FACT SHEET

Baccaurea velutina (Ridley) Ridley

GENUS *Baccaurea* **FAMILY** Phyllanthaceae

COMMON NAMES Tampoi tunggau

EDIBLE PARTS AND USES

Edible parts: Fruit

DESCRIPTION

A tree. It grows 12-35 m tall. The branches have a velvety covering. The leaves are oval and 17 cm long by 10 cm wide. The fruit stalks are 7 cm long. The fruit is like a berry. They are small and orange coloured. They are sour.

HABITAT

A tropical plant. It grows on ridges in the rain forest. It grows up to 1500 m altitude above sea level.

NOTES

There are about 80 Baccaurea species.

REFERENCES

Bircher, A. G. & Bircher, W. H., 2000, Encyclopedia of Fruit Trees and Edible Flowering Plants in Egypt and the Subtropics. AUC Press. p 51

Burkill, I.H., 1966, A Dictionary of the Economic Products of the Malay Peninsula. Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia. Vol 1 (A-H) p 283

Milow, P., et al, 2013, Malaysian species of plants with edible fruits or seeds and their evaluation. International Journal of Fruit Science. 14:1, 1-27

Verheij, E. W. M. and Coronel, R.E., (Eds.), 1991, Plant Resources of South-East Asia. PROSEA No 2. Edible fruits and nuts. Pudoc Wageningen. p 320

World Checklist of Useful Plant Species 2020. Royal Botanic Gardens, Kew

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/9b8315d8-3b36-4b35-a00a-b310bb769d65