PLANT FACT SHEET

Dichelostemma capitatum subsp. pauciflorum (Torr.) Keator

GENUS *Dichelostemma* **FAMILY** Asparagaceae

COMMON NAMES Bluedick

EDIBLE PARTS AND USES

Edible parts: Flowers, Root, Tuber

The corms are eaten raw, fried, roasted or boiled.

DESCRIPTION

A herb that keeps growing from year to year. The bulbs are 2 cm long. The leaves are 5 mm wide and 27-32 cm long. The flowers are 1 cm long and violet to blue.

HABITAT

It grows in wetlands and along rivers. Arizona

NOTES

Possibly now Dichelostemma capitatum ssp. pauciflorum
Also put in the family Themidaceae.

REFERENCES

Beckstrom-Sternberg, Stephen M., and James A. Duke. "The Foodplant Database." http://probe.nalusda.gov:8300/cgi-bin/browse/foodplantdb. (ACEDB version 4.0 - data version July 1994)

Brickell, C. (Ed.), 1999, The Royal Horticultural Society A-Z Encyclopedia of Garden Plants. Convent Garden Books. p 364

CASTETTER & BELL,

CASTETTER & UNDERHILL

Facciola, S., 1998, Cornucopia 2: a Source Book of Edible Plants. Kampong Publications, p 7 (As Brodiaea pulchella)

Hibbert, M., 2002, The Aussie Plant Finder 2002, Florilegium. p 86

Lim, T. K., 2015, Edible Medicinal and Non Medicinal Plants. Volume 9, Modified Stems, Roots, Bulbs. Springer p 33

MINNIS

Plants for a Future database, The Field, Penpol, Lostwithiel, Cornwall, PL22 0NG, UK. http://www.scs.leeds.ac.uk/pfaf/

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/9ccc0c6b-a24f-4232-b0d4-3ab5d344bcd7