

Dioscorea bulbifera

Common name(s)

Potato yam,

Edible portion:

Tuber, Root, Bulbils, Vegetable,



Distribution

A tropical plant. It will grow from the coast up to about 1700 m altitude in equatorial zones. It is common near the edge of grassland and forest at mid altitudes. It grows in savannah woodland. Both wild and cultivated forms occur. It is common in the Philippines, Solomon Islands and Papua New Guinea near secondary forest at low and medium altitudes. It is cultivated in Africa and the West Indies. In Nepal it grows to 2100 m altitude. It suits hardiness zones 9-12. In Yunnan.

Description

A yam with a long smooth stemmed vine, round in cross section. It winds to the left. It does not have spines. The vine can climb up into trees and grow to long lengths. The leaves are large and round. They are pointed at the tip and round at the base. About 7 veins arise from the tip of the leaf stalk. Leaves can be 14-30 cm across and slightly longer than wide. This yam produces bulbils (potatoes) in the angles of the leaves along the vine. These are often flattened and can be grey brown or purple. Under the ground it has a smaller tuber normally covered with roots. The flowers are large. The male flowers are in spikes up to 20 cm long. The female spikes are usually in pairs. The fruit are winged and about 2.5 cm long by 1.5 cm across. The seeds have wings. The bulbils normally have few fibres through the tissue compared to some yams tubers. The flesh of many varieties is yellow.

Use

The tubers are cooked and eaten. More commonly the aerial bulbils are eaten after cooking. Often the cooking water is changed a couple of times to remove bitterness.

Some kinds are bitter and inedible or at least require special processing and cooking.

Some varieties are poisonous. Some can cause act as contraceptives.

Cultivation

Either the bulbils off the vine, or the underground tubers are planted. Because the vines are long, training them up trees is convenient. The bulbils have to be stored for a period of time before they will sprout. The plant is annual and leaves die off for 1-4 months each year before re-sprouting from the tuber. Bulbils only grow shoots from one end unless the bulbil is cut into pieces. If the larger bulbils are cut, the cut surfaces should be dried and healed in a shady place for 2-3 days before planting. Bulbils are planted 8-12 cm below the surface and spacing can be 100 cm by 100 cm or variations of this. Normally nitrogen and potassium fertilisers give greater responses than phosphorus.

Friable well drained soils are most suitable. Often very little cultivation or mounding is done. A high level of organic matter improves yield. Staking is normally required but often trees or living stakes are used. Branched stakes 2 m high are suitable. They need to be strongly erected as vine and tuber growth can be extensive and heavy.

Production

Bulbils or aerial yams are produced as soon as leaves begin to unfold and continue until the plant reaches maturity. These aerial tubers often fall from the plant. Harvesting can start 3 months after planting but immature tubers have less starch. Underground tubers are normally not harvested until the leaves die back. Wounds and damage to the tuber surface normally heals naturally given dry aerated conditions. Some varieties have aerial tubers which are seasonally dormant and only grow after an extended period of storage. Others germinate relatively quickly.

Nutritional values

Edible Part	Moisture %	Energy kJ	Energy kcal	Protein g	Provit A µg	Vit C mg	Iron mg	Zinc mg
(per 100 grams of edible portion)								
Tuber	72.5	455	109	2.3		32	1.1	0.7
Tuber	70.8	357	85	2.7		78	3.1	0.4



Please Note: Except where otherwise noted, content on this site is licensed under a Creative Commons Attribution 3.0 Licence.- this means you can share it freely, as is and with acknowledgement.

Indexing Data (print optional)

Found in

Africa, American Samoa, Andamans, Angola, Asia, Australia, Bangladesh, Benin, Bhutan, Bolivia, Bougainville, Brazil, Burkina Faso, Burundi, Cambodia, Cameroon, Central Africa, Central African Republic, CAR, Central America, China, Christmas Island, Chuuk, Colombia, Comoros, Congo DR, Congo R, Cook Is., Costa Rica, Côte d'Ivoire, Chuuk, Cuba, Dominican Republic, East Africa, East Timor, Ecuador, Ethiopia, Fiji, French Polynesia, FSM, Gambia, Ghana, Guam, Guiana, Guianas, Guinea, Guinée, Guinea-Bissau, Haiti, Himalayas, India, Indochina, Indonesia, Ivory Coast, Jamaica, Japan, Kenya, Korea, Laos, Liberia, Madagascar, Malawi, Malaysia, Maldives, Mali, Marquesas, Mauritius, Micronesia, Myanmar, Namibia, Nepal, New Caledonia, Niger, Nigeria, North America, Northeastern India, Pacific, Pakistan, Palau, Papua New Guinea, PNG, Peru, Philippines, Puerto Rico, Reunion, Rotuma, Sahel, Samoa, Sao Tome and Principe, SE Asia, Senegal, Sierra Leone, Sikkim, Solomon Islands, Southern Africa, South America, Sri Lanka, St Lucia, Suriname, Tahiti, Taiwan, Thailand, Timor-Leste, Togo, Tonga, Tanzania, Uganda, USA, Vanuatu, Venezuela, Vietnam, Wallis & Futuna, West Africa, West Indies, Yap, Zambia, Zimbabwe,

Synonyms

Dioscorea anthropophagorum A. Chev.;
Dioscorea bubifera L. var. *sativa* (Thunb.) Prain;
Dioscorea bulbifera var. *anthropophagorum* (A. Chev.) Summerh.;
Dioscorea latifolia Benth.;
Dioscorea sativa var. *rotunda*;
Dioscorea sativa Thunb.;
Hemia bulbifera (Linnaeus) Kunth.;

Other common names

Abubu, Acom, Aerial Yam, Agbanio, Ah-lu-thi, Air-potato, Akam, Ankindjek, Apueraka, Assidhakattala, Bachi, Ban tarul, Banalu, Bantarul, Ban temee, Basel phauk, Batata de rama, Bayag-toro, Bayal, Bejuco de natilla, Belloi, Bon-alu, Bontarul, Bubaia, Buifu, Cambare marron, Cara de are, Catoco, Chaxo poa, Chedu dumpa, Chedupaddu-dumpa, Cheeky yam, Chitangula, Chupri alu, Coolngarrie, Cu mei, Dai, Danda yam, Dandam, Dandandim-o, Dangkanda, Dau fasia, Dehs prei, Denebra, Dimoa, Drun, Dukkar-kand, Endome, Engin, Favi, Fikengere, Fui, Gaicha alu, Gaihi, Ganmanggu, Gasalu, Gathalu, Geetha, Genebra, Gethi, Gethia kanda, Ghartarul, Gidtha, Githi, Githa, Githo, Goch alu, Gosh alu, Gulgariny, Gunda, Haaran bo, Heggenasum, Helak, Ho, Hofikary, Hoi, Huang du, Huwi blichik, Indrenni, Ignose bois, Ignose-sau-vage, Inga piru, Iroga, Jebubug basu, Kadukand, Kadukaranda, Kadvokand, Kaile, Kaile dranu, Kaile manu, Kalangua, Ka la si, Kalialia, Karanda, Karukarinda, Karuvalli, Kasiena, Kattala, Kattu-kachil, Khoinga, Ki, Kodi, Kodikilangu, Koile mila, Konjo, Kuchung, Kurlama, Lac, Lahara, Lekut, Lindya, Litu, Lu gong lai biu, Mabuia, Magnaheugo, Malakaka yapendalamu, Man nok, Manamund, Manyanya, Mas aloo, Mas alu, Matalu, Matara, Mataru, Methe aloo, Mithene, Mitho githa, Mpenga, Muwana, Name del aire, Name volador, Niambe-de-matom, Nika, Numwe, Ofaka, Ofa lei, Ogoko, Oi, Okoo, Otaheite yam, Pain, Palai, Pannukilangu, Papa aerea, Papa del aire, Papa voladora, Peng-khe, Pesta-alu, Pisimonaju, Piska sanga, Pitaalu, Puralu, Puri, Putsa-u, Ranmataru, Ratalu, Ratulu, Rok, Ruipan, Rukhel, Sang, Shunku papa, Sise, Soi, Soko, Suaralu, Syak, Te'e, Tewe, Tha borok, Thaphu miyung-wablai, Tikor alu, Timbom, Tito githa, To, Ubi atatus, Ubi-ubihan, Udala, Ufi lei, Uhi, Undome, Uwi gantung, Wila, Yi bao, Yoi,

REFERENCES

Potato yam references *Dioscorea bulbifera*

Abbiw, D.K., 1990, Useful Plants of Ghana. West African uses of wild and cultivated plants. Intermediate Technology Publications and the Royal Botanic Gardens, Kew. p 27

Acharya K. P. and Acharya, R., 2010, Eating from the Wild: Indigenous knowledge on wild edible plants in Parroha VDC of Rupandehi District, Central Nepal. International Journal of Social Forestry. 3(1):28-48

Acipa, A. et al, 2013, Nutritional Profile of some Selected Food Plants of Otwal and Ngai Counties, Oyam District, Northern Uganda. African Journal of Food, Agriculture, Nutrition and Development. 13(2)

Agarwal, R. & Chandra, V., 2019, Diversity of wild edible plants in the Mandal-Chopta forest, Uttarakhand. Journal of Medicinal Plants Studies 2019; 7(1): 89-92

Ambasta, S.P. (Ed.), 2000, The Useful Plants of India. CSIR India. p 174

Anderson, E. F., 1993, Plants and people of the Golden Triangle. Dioscorides Press. p 209

Arinathan, V., et al, 2007, Wild edibles used by Palliyars of the western Ghats, Tamil Nadu. Indian Journal of Traditional Knowledge. 6(1) pp 163-168

Arora, K., Indigenous Forest Management in the Andaman and Nicobar Islands, India.

Aryal, K. P. et al, 2009, Uncultivated Plants and Livehood Support - A case study from the Cheopang people of Nepal. Ethnobotany Research and Applications. 7:409-422

Aryal, K. P., et al, 2018, Diversity and use of wild and non-cultivated edible plants in the Western Himalaya. Journal of Ethnobiology and Ethnomedicine (2018) 14:10

Avinash, B. K., Kumar, R., et al, 2013, Ethnobotany of Tharus of Dudhwa National Park, India. Mintage journal of Pharmaceutical & Medical Science 6-11

Awat, T., 1997, A Study on the Ecology and Ethnobotany of Non-cultivated Food Plants and Wild Relatives of Cultivated crops in Gambella Region, Southwestern Ethiopia. Addis Ababa University. p 78

Bandyopadhyay, S. et al, 2009, Wild edible plants of Koch Bihar district, West Bengal. Natural Products Radiance 8(1) 64-72

Bandyopadhyay, S., et al, 2012, A Census of Wild Edible Plants from Howrah District, West Bengal, India. Proceedings of UGC sponsored National Seminar 2012

BARC, 2016, State of Biodiversity for Food and Agriculture in Bangladesh. Bangladesh Agricultural Research Council.

Baro, D., Baruah, S. and Borthukar, S. K. 2015, Documentation on wild vegetables of Baksa district, BTAD (Assam). Scholars Research Library. Archives of Applied Science Research, 2015, 7 (9):19-27

Barrau, J., 1976, Subsistence Agriculture in Polynesia and Micronesia. Bernice P. Bishop Museum, Bulletin 223 Honolulu Hawaii. Kraus reprint. p 45

Barua, U., et al, 2007, Wild edible plants of Majuli island and Darrang districts of Assam. Indian Journal of Traditional Knowledge 6(1) pp 191-194

Bastakoti, R., et al, 2008, Food Insecurity and dependency of Chepang Communities on Wild Edible Plants. Sustainable Forest Management and Poverty Alleviation: Roles of Traditional Forest-related Knowledge IUFRO World Series Volume 21

BHARGAVA,

BHARGAVA, (As *Dioscorea versicolor*)

Bhogaonkar, P. Y. & Devarkar, V. D., 2018, Inventory for ethnovegetable knowledge of the tribals of Satpura hill area-Melghat, Dist. Amravati (India), Plantae Scientia, 2018; Vol. 01 Issue 01. 01 00