



PLANT FACT SHEET

Annona reticulata L.

GENUS *Annona* FAMILY Annonaceae

COMMON NAMES Bullock's heart

EDIBLE PARTS AND USES

Edible parts: Fruit

The ripe fruit is eaten fresh.

They are also used for preserves, drinks, ice cream, custards and other desserts

CAUTION: The seed kernel is poisonous.

DESCRIPTION

A small tree up to 7.5 m tall. It has several branches near the base. Trees lose their leaves at some times of the year. New shoots have short brown hairs but older wood is smooth and shiny. The leaves are long and spear shaped with short leaf stalks. Around the edge of the leaf is a clear edge. The leaves smell when crushed. Flowers are greenish yellow. They occur in groups where the leaves join the stalk. Flowers occur on new wood growth. The fruit are reddish brown in colour and 10-12 cm across. There is a fine hexagonal pattern over the fruit. Inside there are large brown seeds. The fruit are edible but the seeds are not eaten.

HABITAT

A tropical plant. It occurs in the tropical lowlands and grows up to at least 1200 m altitude. In Colombia it grows between 500-1,900 m above sea level. It can grow on poorer soils with different levels of acidity. It cannot stand water-logging. It suits humid climates. It is less suited to dry climates. It can grow in arid places. In Nepal it grows to 900 m altitude. It suits hardiness zones 10-12. In Yunnan. In Brisbane Botanical Gardens.

PRODUCTION

Cultivation

Plants are normally grown from seeds. Seedling trees vary quite a bit. Seedlings are easy to transplant. A spacing of 4-7 m is suitable. Better kinds can be grown using budding or grafting.

Harvesting and Seed Production

Trees begin fruiting at 3-5 years. Fruit setting is often improved by hand pollination. Fruit mature in about 200 days. Fruit can vary from 0.25 to 2.25 kg each. The fruit has 13% sugar. Fruit production is seasonal. A tree may bear 100-150 fruit in a season. The season is normally Dec. to March.

Planting instructions

Bullock's heart

Scientific name: *Annona reticulata*

What is the plant like?

The Bullock's heart is a small tree up to 7.5m tall. It has several branches near the base. New shoots have short brown hairs but older wood is smooth and shiny. Trees lose their leaves at some times of the year.

It has spear shaped leaves. They have short leaf stalks. Around the edge of the leaf is a clear edge. Leaves smell when crushed.

The flowers are greenish yellow. They occur in groups where the leaves join the stalk. They occur on new wood growth.

The fruit are round and about 10 to 12 cm across. They are yellowish brown in colour. Over the surface of the fruit there is a fine hexagonal pattern. There are large brown seeds inside the fruit. The flesh is yellow. The fruit are eaten raw.

Where is the Bullock's heart grown?

The Bullock's heart is a tropical plant. It occurs in the lowlands and up to about 1200 metres altitude. It occurs around some coastal towns.

It is less suited to dry climates.

How do you grow Bullock's heart?

Mostly trees are grown from seed. These trees vary quite a bit. Seedlings are easy to transplant. Trees need to be about 7m apart.

Better kinds can be grown using budding or grafting.

Trees begin to produce fruit after about 3 to 5 years. Trees flower and fruit throughout the year.


NOTES

There are about 100-150 *Annona* species. It has anticancer properties.

PICTURES



Credit: Food Plants International (Bruce French)



Credit: Food Plants International (Bruce French)



Credit: Food Plants International (Bruce French)



Credit: Food Plants International (Bruce French)

REFERENCES

Bullock's heart references

- Ajesh, T. P., et al, 2012, Ethnobotanical Documentation of Wild Edible Fruits used by Muthuvan Tribes of Idukki, Kerala-India. *International Journal of Pharma and Bio Sciences* 3(3): 479-487
- Ambasta S.P. (Ed.), 2000, *The Useful Plants of India*. CSIR India. p 43
- Anderson, E. F., 1993, *Plants and people of the Golden Triangle*. Dioscorides Press. p 202
- Andrare, G. M. M., et al, 2023, Plantas Alimenticias Nao Convencionais (PANC) Nos Levantamentos Etnobotanicos do Brazil. *Novos Cadernos NAEA* v. 25 No. 2 p 198
- Ashton, M. S., et al 1997, *A Field Guide to the Common Trees and Shrubs of Sri Lanka*. WHT Publications Ltd. p 100
- Bajpai, O., et al, 2015, Tree species of the Himalayan Terai region of Uttar Pradesh, India: a checklist. *Check List* 11(4): 1718
- Bandyopadhyay, S. et al, 2009, Wild edible plants of Koch Bihar district, West Bengal. *Natural Products Radiance* 8(1) 64-72
- Bandyopadhyay, S., et al, 2012, A Census of Wild Edible Plants from Howrah District, West Bengal, India. *Proceedings of UGC sponsored National Seminar 2012*
- Banerjee, A., et al, 2013, Ethnobotanical Documentation of Some Wild Edible Plants in Bankura District, West Bengal, India. *The Journal of Ethnobiology and Traditional Medicine*. Photon 120 (2013) 585-590
- Barwick, M., 2004, *Tropical and Subtropical Trees. A Worldwide Encyclopedic Guide*. Thames and Hudson p 27
- Bircher, A. G. & Bircher, W. H., 2000, *Encyclopedia of Fruit Trees and Edible Flowering Plants in Egypt and the Subtropics*. AUC Press. p 31
- Blench, R., 2004, Fruits and Arboriculture in the Indo-Pacific Region. *Indo-Pacific Prehistory Association Bulletin* 24. (Taipei Papers Volume 2) p 34
- Bodkin, F., 1991, *Encyclopedia Botanica*. Cornstalk publishing, p 86
- Borgohain, D., 2017, Ethno-botanical study of wild edible fruits consumed by the people of Lakhimpur district of Assam, India. *Annals of Plant Sciences* 6.11 pp 1728-1735
- Bourret, D., 1981, *Bonnes-Plantes de Nouvelle-Caledonie et des Loyaute*. ORSTOM. p 38
- Brahma, S., et al, 2013, Wild edible fruits of Kokrajhar district of Assam, North-East India, *Asian Journal of Plant Science and Research* 3(6):95-100
- Brouk, B., 1975, *Plants Consumed by Man*. Academic Press, London. p 170
- Brown, 1951, *Useful Plants of the Philippines*.
- Burkill, H. M., 1985, *The useful plants of west tropical Africa*, Vol. 1. Kew.
- Burkill, I. H., 1935, *A Dictionary of the Economic Products of the Malay Peninsula*. p 168
- Casas, A., et al, 1996, Plant Management Among the Nahua and the Mixtec in the Balsas River Basin, Mexico: An Ethnobotanical Approach to the Study of Plant Domestication. *Human Ecology*, Vol. 24, No. 4 pp. 455-478
- Chakraborty, S. & Chaturbedi, H. P., 2014, Some Wild Edible Fruits of Tripura- A Survey. *Indian Journal of Applied research*. (4) 9
- Chandrashekar, U. M., 2009, Tree species yielding edible fruit in the coffee-based homegardens of Kerala, India: their diversity, uses and management. *Food Sec.* 1:361-370
- Chiang Mai University Vol. 4 , 1964, Table 6 p 45
- Chin, H.F., & Yong, H.S., 1996, *Malaysian Fruits in Colour*. Tropical press, Kuala Lumpur p 57
- Chizmar Fernandez, C., et al, 2009, Plantas comestibles de Centroamerica. Instituto de Biodiversidad, Costa Rica. p 38
- Chowdhury, T., et al, 2014, Wild edible plants of Uttar Dinajpur District, West Bengal. *Life Science Leaflets*. 47:pp 20-36
<http://lifesciencesleaflets.ning.com>
- Chowdhury, M. & Mukherjee, R., 2012, Wild Edible Plants Consumed by Local Communities of Maldah of West Bengal, India. *Indian J.Sci.Res.*3(2) : 163-170
- Cooper, W. and Cooper, W., 2004, *Fruits of the Australian Tropical Rainforest*. Nokomis Editions, Victoria, Australia. p 19
- Coronel, R.E., 1982, *Fruit Collections in the Philippines*. IBPGR Newsletter p 6
- Cosiaux, A., 2014, Local uses of tree species and contribution of mixed tree gardens in Saleman. Village near Manusela National Park, Seram Island, Maluku (Indonesia) CIFOR working paper 137. p 38
- Dahlgren, B. E. & Standley, P. C., 1944, *Edible and Poisonous plants of the Caribbean Region*. Dept. of the Navy. p 22
- Darley, J.J., 1993, *Know and Enjoy Tropical Fruit*. P & S Publishers. p 2
- Das, S. and De, B., 2013, Evaluation of Angiotension I-Converting Enzyme (ACE) inhibitory potential of some underutilized indigenous fruits of West Bengal using an in vitro model. *Fruits*, Vol. 68:499-506
- Das, T. & Das, A. K., 2005, Inventorying plant biodiversity in homegardens: A case study in Barak Valley, Assam, North East India. *CURRENT SCIENCE*, VOL. 89, NO. 1, 10 JULY 2005
- Datar, M. N. & Upadhye, A. S., 2015, Forest foods of Northern Western Ghats: Mode of Consumption, Nutrition, and Availability. *Asian Agri-History* Vol. 19, No. 4, 2015 (293-316)
- Dobriyal, M. J. R. & Dobriyal, R., 2014, Non Wood Forest Produce an Option for Ethnic Food and Nutritional Security in India. *Int. J. of Usuf. Mngt.* 15(1):17-37
- Duarte, M. C., et al, 2022, Diversity of Useful Plants in Cabo Verde Islands: A Biogeographic and Conservation Perspective. *Plants* 2022, 11, 1313 p 8

Full reference list (105 more) at <https://www.echocommunity.org/en/resources/b012eb07-a973-449d-9b56-b774868c6e31>

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE [CC BY-NC-SA 3.0](#)

ONLINE FACT SHEET <https://www.echocommunity.org/en/resources/b012eb07-a973-449d-9b56-b774868c6e31>