PLANT FACT SHEET

Santiria laevigata Blume

GENUS *Santiria* **FAMILY** Burseraceae

COMMON NAMES Kedongdong kerut

EDIBLE PARTS AND USES

Edible parts: Fruit

DESCRIPTION

A large tree. It grows 30 m tall and the trunk is 80 cm across. It has buttresses 4 m high. The bark is grey brown and cracked and scaly. The leaves have 1-4 pairs of leaflets. They are 7-12 cm long by 3-7 cm wide. They are rounded at the base and have a pointed tip. There are 7-15 pairs of side veins. They are raised on both surfaces. The flowers are in groups in the axils of leaves. The fruit can be rounded or oblong. They are 1-2 cm across.

HABITAT

A tropical plant. It can grow on poor and swampy soils from sea level to 1,200 m above sea level.

REFERENCES

Altschul, S.V.R., 1973, Drugs and Foods from Little-known Plants. Notes in Harvard University Herbaria. Harvard Univ. Press. Massachusetts. no. 1990

Burkill, I.H., 1966, A Dictionary of the Economic Products of the Malay Peninsula. Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia. Vol 2 (I-Z) p 1991

Leenhouts, P.W., 1955, Burseraceae in Flora Malesiana Vol 5 p 232

Slik, F., www.asianplant.net

Soepadmo, E. and Wong, K. M., 1995, Tree Flora of Sabah and Sarawak. Forestry Malaysia. Volume One. p 90

Suwardi, A. B., et al, 2020, Wild edible fruits generate substantial income for local people of the Gunung Leuser National Park, Aceh Tamiang Region. Ethnobotany Research & Applications 20:11

Suwardi, A.B., et al, 2023, The diversity and regional conservation status of wild edible fruit species in Sumatra, Indonesia. Biodiversitas Vol. 24 No. 6. pp 3245-3257

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/c6c77854-aa00-439d-9cf1-30c2b5a3067b