PLANT FACT SHEET

Hypholoma sublateritium (Fr.) Quel.

GENUS *Hypholoma* **FAMILY** Strophariaceae

COMMON NAMES Brick Cap, Chestnut mushroom, Perplexing hypholoma

EDIBLE PARTS AND USES

Edible parts: Fungus, Mushroom

Caution; It is normally very bitter and not edible.
CAUTION: It is poisonous.
The fruiting bodies are boiled and eaten or used in soups. They are normally treated in vinegar before cooking.
Cultivated forms are available.

DESCRIPTION

A mushroom. The young caps are brick red. The veil remnants hang on the edge. They have yellowish-green gills. At maturity the cap fades to orange-brown and the gills become purple-black due to spore production.

HABITAT

It is a temperate plant.

REFERENCES

Facciola, S., 1998, Cornucopia 2: a Source Book of Edible Plants. Kampong Publications, p 253

Fuhrer, B & Robinson, R., 1992, Rainforest Fungi of Tasmania and SE Australia, CSIRO p 37

Fuhrer, B., 2005, A field guide to Australian Fungi. Bloomings Books. p 97

Hwang, L. C., et al, 2020, Traditional Botanical Uses of Non-Timber Forest Products (NTFP) in Seven Counties in Liberia. ACS Symposium Series; American Chemical Society: Washington, DC, 2020.

Jordan, P., 2000, The Mushroom Guide and Identifier, Hermes House, p 60

Pace, G., 1998, Mushrooms of the world. Firefly books. p 56 (Not edible)

Tanaka,

Uphof,

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/c97bcfee-9b07-4637-b2d1-cff598b97a0d