PLANT FACT SHEET

Bromelia hieronymi Mez

GENUS *Bromelia* **FAMILY** Bromeliaceae

COMMON NAMES Kisye, Isa

EDIBLE PARTS AND USES

Edible parts: Bud, Stolons, Stem tips

The very young stem buds are eaten with fish fats.
The young stolons are peeled and eaten with fish fats.
The heart of the plant is roasted on the fire.

DESCRIPTION

A pineapple like herb. It has stolons. The leaves are in a ring. The fruit form fleshy berries.

HABITAT

It is a subtropical plant. In Argentina it grows between 400-700 m above sea level. It is usually in seasonally dry forests.

REFERENCES

Arenas, P. and Scarpa, G. F., 2006, Edible wild plants of the Chorote Indians, Gran Chaco, Argentina. Botanical Journal of the Linnean Society. Vol. 153 (1): pp 73-85

Kermath, B. M., et al, 2014, Food Plants in the Americas: A survey of the domesticated, cultivated and wild plants used for Human food in North, Central and South America and the Caribbean. On line draft. p 145

Montani, M. C. & Scarpa, G. F., 2016, Recursos vegetales y prácticas alimentarias entre indígenas tapiete del noreste de la provincia de Salta, Argentina. Darwiniana, nueva serie vol.4 no.1 San Isidro jul. 2016

Polini, G., et al, Useful and edible plants of Paraguay Region of Chaco. p 26

Scarpa, G. F., 2009, Wild food plants used by the indigenous peoples of South American Gran Chaco: A general synopsis and intercultural comparison. Journal of Applied Botany and Food Quality 83:90-101

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/ed116855-e406-4711-96fd-bd62807c03bf