PLANT FACT SHEET

Corymbia dampieri (D.J. Carr & S.G.M. Carr) K.D. Hill & L.A.S. Johnson

GENUS *Corymbia* **FAMILY** Myrtaceae

COMMON NAMES Dampier's Bloodwood

EDIBLE PARTS AND USES

Edible parts: Gall

DESCRIPTION

A tree. It grows 5-11 m high. The bark is rough and cracked into sections. The flowers are cream and white.

HABITAT

A tropical plant. It grows from tropical to temperate regions in Western Australia. It grows on red sand or clay soils.

NOTES

There are a 110 Corymbia species mostly originally in Australia.

REFERENCES

Kenneally, K.E., Edinger, D. C., and Willing T., 1996, Broome and Beyond, Plants and People of the Dampier Peninsula, Kimberley, Western Australia. Department of Conservation and Land Management. p 142

Paczkowska, G . & Chapman, A.R., 2000, The Western Australian Flora. A Descriptive Catalogue. Western Australian Herbarium. p 357

Wheeler, J.R.(ed.), 1992, Flora of the Kimberley Region. CALM, Western Australian Herbarium, p 517

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/ef0b7f8e-71b3-4cd8-a054-754147b992e3