PLANT FACT SHEET

Araujia brachystephana (Griseb.) Fontella & Goyder

GENUS *Araujia* **FAMILY** Apocynaceae

EDIBLE PARTS AND USES

Edible parts: Fruit, Latex

The fruit is eaten fresh and boiled.
The latex of the branch is used to curdle milk.

DESCRIPTION

A creeper or vine.

HABITAT

It is a subtropical plant.

NOTES

There were 10 Morrenia species - now Araujia. They occur in South America. They are also put in the family Asclepiadaceae.

REFERENCES

Arenas, P, 1999, Morrenia odorata (Asclepiadaceae), an Edible Plant of the Gran Chaco. Economic Botany Vol. 53. No. 1. pp. 89-97 (As Morrenia brachystephana)

Palmieri, V. S., et al, 2018, Aproximaciones etnobotánicas de las especies y prácticas de frutos nativos comestibles de la Actualidad. Aportes para la interpretación del pasado prehispánico de cerro colorado (Córdoba, Argentina). Bol. Soc. Argent. Bot. 53 (1): 115-133

Palmeiri, V. S., & Geisa, M. G., 2019, Las plantas comestibles empleadas por las comunidades comechingonas de San Marcos Sierras (Córdoba, Argentina). Primeras aproximaciones. Bol. Soc. Argent. Bot. 54: 295-309.

Palmieri, V. S., et al, 2022, Wild edible plants of the Central Mountains in Argentina. Comparing subregions to understand the complexity of local botanical knowledge. Rodriguésia 73: e01092021. 2022

Polini, G., et al, Useful and edible plants of Paraguay Region of Chaco. p 12

Scarpa, G. F., 2009, Wild food plants used by the indigenous peoples of South American Gran Chaco: A general synopsis and intercultural comparison. Journal of Applied Botany and Food Quality 83:90-101 (As Morrenia brachystephana)

PROVIDED BY **Food Plants International**

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET https://www.echocommunity.org/en/resources/f548069d-812f-4054-8c10-c90ec2ec1b13