



PLANT FACT SHEET

Carpotroche longifolia (Poepp.) Benth.

GENUS *Carpotroche* FAMILY Achariaceae

COMMON NAMES Longleaf carpotroche

EDIBLE PARTS AND USES

Edible parts: seeds - aril, Fruit, Seeds

The fruit are eaten. It is the small fleshy layer (aril) around the seeds that is eaten.

Caution: The fruit have a cyanide smell and may need cooking.

DESCRIPTION

A shrub or tree. It grows 12 m tall. The flowers are in small clusters. They grow on the old wood. The fruit is a narrowly oval, woody capsule. It is 5 cm long by 3 cm wide and have 10-12 wings. These are a light green with small teeth. There are wings between these also with teeth. There are about 50 seeds with one rounded and 2 flat sides. They are 1 cm long by 0.6 cm wide and with red or orange arils or coverings over the seeds.

HABITAT

A tropical plant. It grows along creeks and rivers, expecially on slopes.

PRODUCTION

Harvesting and Seed Production

Fruit are harvested off the tree trunk.

NOTES

There are about 30 *Carpotroche* species. They are also put in the family Kiggelariaceae and Euphorbiaceae. Also put in the Flacourtiaceae family.

REFERENCES

- Bircher, A. G. & Bircher, W. H., 2000, Encyclopedia of Fruit Trees and Edible Flowering Plants in Egypt and the Subtropics. AUC Press. p 85
- Cuatrec. Publication : Trop. Woods No. 101, 28 (1955)
Basionym author: (R.E.Schult.) Publishing author: Benth. Publication: in Journ. Linn. Soc. v. Suppl. ii. 82 1861
- Etkin, N.L. (Ed.), 1994, Eating on the Wild Side, Univ. of Arizona. p 137 (As *Carpotroche linguifolia*)
- Grandtner, M. M., 2008, World Dictionary of Trees. Wood and Forest Science Department. Laval University, Quebec, Qc Canada. (Internet database <http://www.WDT.QC.ca>)
- Grandtner, M. M. & Chevette, J., 2013, Dictionary of Trees, Volume 2: South America: Nomenclature, Taxonomy and Ecology. Academic Press p 102
- Kermath, B. M., et al, 2014, Food Plants in the Americas: A survey of the domesticated, cultivated and wild plants used for Human food in North, Central and South America and the Caribbean. On line draft. p 187
- Penafiel Anchundia, D. D., 2017, Traditional food consumption and its nutritional contribution in Guasaganda, Central Ecuador. PhD Ghent University p 71

Roa, J. A. G. & Boada, D. S. G., 2018, Fundación para el Fortalecimiento de la Fruticultura y Plantas Alimenticias no Convencionales en Colombia.

van Roosmalen, M.G.M., 1985, Fruits of the Guianan Flora. Utrecht Univ. & Wageningen Univ. p 126

Vasquez, R. and Gentry, A. H., 1989, Use and Misuse of Forest-harvested Fruits in the Iquitos Area. Conservation Biology 3(4): 350f

PROVIDED BY Food Plants International

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET <https://www.echocommunity.org/en/resources/f6afb9d9-14de-4bbb-a772-600133cc889a>