



PLANT FACT SHEET

Cratoxylum formosum (Jack) Benth. & Hook. f. ex Dyer

GENUS *Cratoxylum* **FAMILY** Hypericaceae

COMMON NAMES Pink mempat

EDIBLE PARTS AND USES

Edible parts: Leaves, bud, Tea

Young leaves are eaten raw or boiled with fish or pork. They give a sour taste. They are used as a fresh vegetable and also in soup.

DESCRIPTION

A shrub or small tree. It grows 1-8 m high. The trunk has long thorns on the lower parts. The bark peels off in flakes. The leaves are simple and opposite. They are lighter green underneath. They are oblong and 4-10 cm long by 2-4 cm wide. The flowers are in groups of 5-8 in the axils of the fallen leaves. The flowers are 1.3 cm across. The fruit is a dark brown capsule that is oblong and 1-1.5 cm long. There are 6-8 seeds per section.

HABITAT


A tropical plant. It grows in clear forests on sandy soils. It grows in dry mixed forests up to 1,000 m above sea level. In XTBG Yunnan.

PRODUCTION

Cultivation

Plants can be grown from seeds or hardwood cuttings.

PICTURES



Credit: Food Plants International (Bruce French)

REFERENCES

- Bhatt, D., Chandra Sekar, K. and Kumar, K., 2020, Tree Diversity of Indian Himalayan Region. G.B. Pant National Institute of Himalayan Environment, Almora, Uttarakhand, India. p 61
- Carig, Elizabeth T., 2020, Guidebook on Native Trees within the Quirino Forest Landscape. Plaridel, Bulacan, St. Andrew Publishing House. p 103
- Cengel, D. J. & Dany. C., (Eds), 2016, Integrating Forest Biodiversity Resource Management and Sustainable Community Livelihood Development in the Preah Vihear Protected Forest. International Tropical Timber Organization p 110
- Cruz-Garcia, G. S., & Price, L. L., 2011, Ethnobotanical investigation of 'wild' food plants used by rice farmers in Kalasin, Northeast Thailand. *Journal of Ethnobiology and Ethnomedicine* 7:33
- Eiadthong, W., et al, 2010, Management of the Emerald Triangle Protected Forests Complex. Botanical Consultant Technical Report. p 23
- Fu, Yongneng, et al, 2003, Relocating Plants from Swidden Fallows to Gardens in Southwestern China. *Economic Botany*, 57(3): 389-402
- Japanese International Research Centre for Agricultural Science www.jircas.affrc.go.jp/project/value_addition/Vegetables
- Kachenchart, B., et al, 2008, Phenology of Edible Plants at Sakaerat Forest. In *Proceedings of the FORTROP II: Tropical Forestry Change in a Changing World*. Bangkok, Thailand.
- Khumgratok, S., Edible Plants in Cultural Forests of Northeastern Thailand. Mahasarakham University Thailand.
- Maituthisakul, P., et al, 2007, Assessment of phenolic content and free radical-scavenging capacity of some Thai indigenous plants. *Food Chemistry* 100: 1409-1418
- Nakahara, K. et al, 2002, Antimutagenicity of Some Edible Thai Plants, and a Biocative Carbazole Alkaloid, Mahanine, Isolated from *Micromelum minutum*. *Journal of Agricultural and Food Chemistry*. 50: 4796-4892
- Nguyen, C. H., et al, 2021, Edible wild plants in the flora of Pu Luong Nature Reserve, Thanh Hoa Province, northern Vietnam. *IOP Conf. Ser.: Earth Environ. Sci.* 876
- Phatlamphu, N., et al, 2021, Ethnobotany of edible plants in Muang District, Kalasin Province, Thailand, *Biodiversitas* Vol. 22, No. 12 pp 5432-5444
- Phon, P., 2000, Plants used in Cambodia. © Pauline Dy Phon, Phnom Penh, Cambodia. p 177
- Slik, F., www.asianplant.net
- Somnasang, P., Moreno, G and Chusil K., 1998, Indigenous knowledge of wild hunting and gathering in north-east Thailand. *Food and Nutrition Bulletin* 19(4) p 359f
- Somnasang, P., et al, 2000, Knowing gathering and eating: Knowledge and attitudes about wild food in an Isan Village in north eastern Thailand. *Journal of Ethnobiology* 20(2):197-216
- Srichaiwong, P., et al, 2014, A Study of the Biodiversity of Natural Food Production to Support Community Upstream of Chi Basin, Thailand. *Asian Social Science* 10 (2):
- Suksri, S., et al, 2005, Ethnobotany in Bung Khong Long Non-Hunting Area, Northeast Thailand. *Kasetsart J., (Nat. Sci)* 39: 519-533
- Sukarya, D. G., (Ed.) 2013, 3,500 Plant Species of the Botanic Gardens of Indonesia. LIPI p 211
- Suwardi, A.B., et al, 2023, The diversity and regional conservation status of wild edible fruit species in Sumatra, Indonesia. *Biodiversitas* Vol. 24 No. 6. pp 3245-3257
- Thitiprasert, W., et al, 2007, Country report on the State of Plant Genetic Resources for Food and Agriculture in Thailand (1997-2004). FAO p 95
- Thorng, R., et al, 2015, Roles of Wild Food Plants in Ethnic Group Communities in Mondulkiri Province, Northeastern Cambodia. *Journal of Mekong Societies* Vol.11 No.2 May-August 2015 pp. 1-17
- Turreira Garcia, N., et al, 2017, Ethnobotanical knowledge of the Kuy and Khmer people in Prey Lang, Cambodia. *Cambodian Journal of Natural History* 2017 (1): 76-101
- Waiyaput, W., et al, 2012, Inhibitory effects of crude extracts from some edible Thai plants against replication of hepatitis B virus and human liver cancer cells. *BMC Complementary and Alternative Medicine*. 12:246

PROVIDED BY Food Plants International

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET <https://www.echocommunity.org/en/resources/f6da3a20-767e-416b-b0b9-26e2982c7d3a>