



PLANT FACT SHEET

Crotalaria ochroleuca G. Don

GENUS *Crotalaria* FAMILY Fabaceae

EDIBLE PARTS AND USES

Edible parts: Leaves, Vegetable, Flowers

The leaves are cooked and eaten. The leaves are often dried. They are also chopped and boiled and eaten with other foods.

Caution: Drying often is used to get rid of a poison.

People become drowsy if they eat the flowers. Dried flowers are used for flavouring.

DESCRIPTION

An erect annual, or short lived perennial. It grows 0.5-2.7 m tall. There are fine hairs on the stems. The leaves have 3 leaflets. These are narrow and 5-13 cm long by 0.5-3 cm wide. They are slightly hairy underneath. The flower clusters are 10-50 cm long, with many flowers which are 2 cm long. They are yellow with red streaks. The wings of the flowers have a purple mark at the base. The fruit is like a cylinder and is 5-7 cm long by 1.5-2 cm wide.

HABITAT

A tropical plant. In Papua New Guinea it grows from 30 m to 1,800 m altitude. It is mostly on plantations. In Zambia it grows in savannah. In southern China it grows in shady places with thin soil.

PRODUCTION

Cultivation

Plants can be grown from seeds.

Harvesting and Seed Production

Young leaves are picked off the plant.

NOTES

There are about 550 *Crotalaria* species. They are mostly tropical.

REFERENCES

Abbiw, D.K., 1990, Useful Plants of Ghana. West African uses of wild and cultivated plants. Intermediate Technology Publications and the Royal Botanic Gardens, Kew. p 42

Abukutsa-Onyango, M.O., 2004. *Crotalaria brevidens* Benth. [Internet] Record from Protabase. Grubben, G.J.H. & Denton, O.A. (Editors). PROTA (Plant Resources of Tropical Africa), Wageningen, Netherlands. < <http://database.prota.org/search.htm>>. Accessed 15 October 2009.

- Abukutsa-Onyango, Mary, 2007, The diversity of cultivated African leafy vegetables in three communities in western Kenya. *AJFAND*, Volume 7, No. 3
- Acipa, A. et al, 2013, Nutritional Profile of some Selected Food Plants of Otwal and Ngai Counties, Oyam District, Northern Uganda. *African Journal of Food, Agriculture, Nutrition and Development*. 13(2)
- Agea, J. G., et al 2011, Wild and Semi-wild Food Plants of Bunyoro-Kitara Kingdom of Uganda: etc. *Environmental Research Journal* 5(2) 74-86
- Andabati, B., & Muyonga, J., 2014, Phenolic content and antioxidant activity of selected Ugandan traditional medicinal foods. *African Journal of Food Science*. 8(8), pp 427-434
- Burkill, H. M., 1985, The useful plants of west tropical Africa, Vol. 3. Kew. Food Composition Tables for use in Africa FAO <http://www.fao.org/infoods/directory> No. 604
- Fowler, D. G., 2007, *Zambian Plants: Their Vernacular Names and Uses*. Kew. p 80
- Gen. hist. 2:138. 1832
- Godfrey, J. et al, 2013, Harvesting, preparation and preservation of commonly consumed wild and semi-wild food plants in Bunyoro-Kitara Kingdom, Uganda. *Int. J. Med. Arom. Plants*. Vol.3 No.2 pp 262-282
- Goode, P., 1989, *Edible Plants of Uganda*. FAO p 39
- Grubben, G. J. H. and Denton, O. A. (eds), 2004, *Plant Resources of Tropical Africa 2. Vegetables*. PROTA, Wageningen, Netherlands. p 230
- Harris, D. J., 2002, The vascular plants of the Dzanga-Sangha Reserve, Central African Republic. *National Botanic Garden of Belgium*, 2002. - 274 pages p 116
- INFOODSUpdatedFGU-list.xls
- Jardin, C., 1970, List of Foods Used In Africa, FAO Nutrition Information Document Series No 2.p 71
- Malaisse, F., 1997, *Se nourrir en floret claire africaine. Approche ecologique et nutritionnelle*. CTA., p 59
- Martin, F.W. & Ruberte, R.M., 1979, *Edible Leaves of the Tropics*. Antillian College Press, Mayaguez, Puerto Rico. p 198
- Masters, T., 2021, Traditional food plants of the upper Aswa River catchment of northern Uganda—a cultural crossroads. *Journal of Ethnobiology and Ethnomedicine* (2021) 17:24
- Maundu, P. et al, 1999, *Traditional Food Plants of Kenya*. National Museum of Kenya. 288p
- Mutie, F. M., et al, 2023, Important Medicinal and Food Taxa (Orders and Families) in Kenya, Based on Three Quantitative Approaches. *Plants* 2023, 12, 1145
- Nyero, A., et al, 2021, Wild and semi-wild edible plants used by the communities of Acholi sub-region, Northern Uganda. *Ethnobotany Research & Applications* 21:16
- Peters, C. R., O'Brien, E. M., and Drummond, R.B., 1992, *Edible Wild plants of Sub-saharan Africa*. Kew. p 132
- Schippers, R.R., 2004. *Crotalaria ochroleuca* G.Don. [Internet] Record from Protabase. Grubben, G.J.H. & Denton, O.A. (Editors). PROTA (Plant Resources of Tropical Africa), Wageningen, Netherlands. < <http://database.prota.org/search.htm>>. Accessed 15 October 2009
- Swaziland's Flora Database <http://www.sntc.org.sz/flora>
- Termote, C., et al, 2014, Assessing the potential of wild foods to reduce the cost of a nutritionally adequate diet: An example from eastern Baringo District, Kenya. *Food and Nutrition Bulletin*, vol. 35, no. 4
- Verdcourt, B., 1979, *Manual of New Guinea Legumes*. Botany Bulletin No 11, Division of Botany, Lae, Papua New Guinea. p 582
- Weinberger, K., Msuya J., 2004, *Indigenous Vegetables in Tanzania. Significance and Prospects*. Technical Bulletin No. 31 AVRDC, Taiwan/FAO p 4
- Williamson, J., 2005, *Useful Plants of Malawi*. 3rd. Edition. Mdadzi Book Trust. p 83
- World Checklist of Useful Plant Species 2020. Royal Botanic Gardens, Kew

PROVIDED BY Food Plants International

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET <https://www.echocommunity.org/en/resources/f966037a-8b30-4b85-8583-e30a3505ebaf>