



PLANT FACT SHEET

Sideroxylon lanuginosum Michx.

GENUS *Sideroxylon* FAMILY Sapotaceae

COMMON NAMES Woolly buckthorn

EDIBLE PARTS AND USES

Edible parts: Fruit, Gum, Sap

The gum is chewed.

The fruit can be eaten. The fruit are ground and mixed with water as a soft drink.

CAUTION: If the fruit are eaten in a large quantities they can cause dizziness and stomach upsets.

DESCRIPTION

A tree. It grows 18 m high. It is usually smaller in cultivation. It loses its leaves in cooler regions but stays evergreen in warmer places. It has a tall narrow crown. The bark is scaly and dark brown. The leaves are small and blunt tipped. They are shiny dark green on the upper surface and are densely woolly underneath. The flowers are 3 mm wide. They are bell shaped and have a 5-lobed white corolla. The fruit are purplish black. They are 12-25 mm long.

HABITAT

It is a temperate and subtropical plant. It grows in valleys and rocky slopes. It usually grows to 760 m altitude in SE of the USA. It suits hardiness zones 6-11.

NOTES

There are about 25 Bumelia species. The Bumelia may be merged with the Sideroxylon.

REFERENCES

- Beckstrom-Sternberg, Stephen M., and James A. Duke. "The Foodplant Database."
<http://probe.nalusda.gov:8300/cgi-bin/browse/foodplantdb>.
(ACEDB version 4.0 - data version July 1994) (As Bumelia lanuginosa)
- Bircher, A. G. & Bircher, W. H., 2000, Encyclopedia of Fruit Trees and Edible Flowering Plants in Egypt and the Subtropics. AUC Press. p 68 (As Bumelia lanuginosa)
- Etherington, K., & Imwold, D., (Eds), 2001, Botanica's Trees & Shrubs. The illustrated A-Z of over 8500 trees and shrubs. Random House, Australia. p 148 (As Bumelia lanuginosa)
- Estrada-Castillon, E., et al, 2014, Ethnobotany in Rayones, Nuevo Leon, Mexico. Journal of Ethnobiology and Ethnomedicine, 10:62
- Grandtner, M. M., 2008, World Dictionary of Trees. Wood and Forest Science Department. Laval University, Quebec, Qc Canada. (Internet database <http://www.wdt.qc.ca>)
- Hedrick, U.P., 1919, (Ed.), Sturtevant's edible plants of the world. p 138 (As Bumelia lanuginosa)
- <http://www.botanic-gardens-ljubljana.com/en/plants>
- Kermath, B. M., et al, 2014, Food Plants in the Americas: A survey of the domesticated, cultivated and wild plants used for Human food in North, Central and South America and the Caribbean. On line draft. p 803
- Little, E.L., 1980, National Audubon Society Field Guide to North American Trees. Alfred A. Knopf. p 630 (As Bumelia lanuginosa)
- Moerman, D. F., 2010, Native American Ethnobotany. Timber Press. p 530

Plants for a Future database, The Field, Penpol, Lostwithiel, Cornwall,
PL22 0NG, UK.
<http://www.scs.leeds.ac.uk/pfaf/> (As Bumelia lanuginosa)

Syn. pl. 1:237. 1805 (As Bumelia lanuginosa)

PROVIDED BY Food Plants International

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE [CC BY-NC-SA 3.0](#)

ONLINE FACT SHEET <https://www.echocommunity.org/en/resources/f96850e9-a7b3-44e1-98d0-44d4770a4ac2>