



PLANT FACT SHEET

Grewia flava DC.

GENUS *Grewia* FAMILY Malvaceae

COMMON NAMES Brandy bush, Raisin bush

EDIBLE PARTS AND USES

Edible parts: Fruit, Leaves, Gum, Roots - coffee

The fruit are eaten either fresh or after being pulped in a mortar. They are made into porridge. They are used for jam and juice.

They are also used to make brandy and beer.

The fruit are dried and eaten with dried locusts. The sun dried fruit can be stored.

DESCRIPTION

A shrub which loses its leaves during the year. It is spreading, branched and twiggy. It can grow up to 4 m tall. Young branches are grey and hairy. The leaves are pale green and rough. They are 1.3-8 cm long and 2.5 cm wide. The tips are rounded and the base is narrowed. The base is unequal sided. The edges have fine teeth. The flowers are small and star shaped. They have narrow yellow petals. They are usually produced singly in the axils of leaves. The fruit have 1 or 2 lobes. They are hairy and red when ripe. They are edible.

HABITAT

A tropical plant. It grows in dry bush regions. It is often on sandy or limestone soils. It is resistant to drought and frost. It grows in area with an annual rainfall between 50-600 mm. It grows between 150-1,830 m above sea level. It can grow in arid places.

PRODUCTION

Cultivation

Plants can be grown from seeds. Seeds must be cleaned and dried in the shade. They are soaked in hot water and left overnight. Seeds do not germinate easily. They take about 24 days to germinate. Seedlings should be transplanted at the 2 leaf stage.

Harvesting and Seed Production

It is fast growing.

NOTES

Fruit are high in sugar. There are about 200 *Grewia* species. They are mostly tropical. The fruit of most may be edible. The dried fruit are about 64% sugar and 4 % protein. These were in the Sparrmanniaceae and the Tiliaceae.

REFERENCES

- Cat. pl. horti monsp. 113. 1813
- Exell, A.W. et al, (Ed), 1963, Flora Zambesiaca Vol 2 Part 1 Crown Agents, London. p 52
- FAO, 1988, Traditional Food Plants, FAO Food and Nutrition Paper 42. FAO Rome p 307
- Fox, F. W. & Young, M. E. N., 1982, Food from the Veld. Delta Books. p 351
- Grivetti, 1976, 1979,
- Grivetti, L. E., 1980, Agricultural development: present and potential role of edible wild plants. Part 2: Sub-Saharan Africa, Report to the Department of State Agency for International Development. p 79
- Heinz & Maguire, 1974,
- INFOODS:FAO/INFOODS Databases
- Jardin, C., 1970, List of Foods Used In Africa, FAO Nutrition Information Document Series No 2.p 140
- Joffe, P., 2007, Creative Gardening with Indigenous Plants. A South African Guide. Briza. p 190
- Lee, 1979,
- Leger, S., 1997, A Description of Today's Use of Plants in West Bushmanland (Namibia). German Development Service. PO Box 220035, 14061 Berlin, Germany. <http://www.sigridleger.de/book/>
- Long, C., 2005, Swaziland's Flora - siSwati names and Uses <http://www.sntc.org.sz/flora/>
- Maguire, 1978,
- Malan & Owen-Smith, 1974,
- Mannheimer, C. A. & Curtis. B.A. (eds), 2009, Le Roux and Muller's Field Guide to the Trees and Shrubs of Namibia. Windhoek: Macmillan Education Namibia. p 318
- Marshall, 1976,
- Mothanka, D. M. T., et al, 2008, Edible Indigenous Fruit Plants of Eastern Botswana. International Journal of Poultry Science. 7(5): 457-460
- Neelo, J., et al, 2015, Ethnobotanical Survey of Woody Plants in Shorobe and Xobe Villages, Northwest Region of Botswana. Ethnobotany Research & Applications 14:367-379
- Neudeck, L. et al, 2012, The Contribution of Edible Wild Plants to Food Security, Dietary Diversity and Income of Households in Shorobe Village, Northern Botswana. Ethnobotany Research & Applications 10:449-462
- Palgrave, K.C., 1996, Trees of Southern Africa. Struik Publishers. p 570
- Palmer, E and Pitman, N., 1972, Trees of Southern Africa. Vol. 2. A.A. Balkema, Cape Town p 1438
- Peters, C. R., O'Brien, E. M., and Drummond, R.B., 1992, Edible Wild plants of Sub-saharan Africa. Kew. p 191
- Plowes, N. J. & Taylor, F. W., 1997, The Processing of Indigenous Fruits and other Wildfoods of Southern Africa. in Smartt, L. & Haq. (Eds) Domestication, Production and Utilization of New Crops. ICUC p 191
- Rodin, 1985,
- Roodt, V., 1998, Trees & Shrubs of the Okavango Delta. Medicinal Uses and Nutritional value. The Shell Field Guide Series: Part 1. Shell Botswana. p 77
- Royal Botanic Gardens, Kew (1999). Survey of Economic Plants for Arid and Semi-Arid Lands (SEPASAL) database. Published on the Internet; <http://www.rbgekew.org.uk/ceb/sepasal/internet> [Accessed 11th June 2011]
- Ruiters-Welcome, A. K., 2019, Food plants of southern Africa. Ph.D. thesis. Univ. of Johannesburg p 75
- Schmidt, E., Lotter, M., & McClelland, W., 2007, Trees and shrubs of Mpumalanga and Kruger National Park. Jacana Media p 402
- Silberbauer, 1965, 1981,
- Story, 1958,
- Swaziland's Flora Database <http://www.sntc.org.sz/flora>
- Tanaka, 1980,
- Tredgold, M.H., 1986, Food Plants of Zimbabwe. Mambo Press. p 70
- van Wyk, Be., & Gericke, N., 2007, People's plants. A Guide to Useful Plants of Southern Africa. Briza. p 44
- van Wyk, B, van Wyk, P, and van Wyk B., 2000, Photographic guide to Trees of Southern Africa. Briza. p 10, 165
- van Wyk, B-E., 2011, The potential of South African plants in the development of new food and beverage products. South African Journal of Botany 77 (2011) 857-868
- Venter, F & J., 2009, Making the most of Indigenous Trees. Briza. p 180
- Wehmeyer, A. S, 1986, Edible Wild Plants of Southern Africa. Data on the Nutrient Contents of over 300 species
- Welcome, A. K. & Van Wyk, B.-E., 2019, An inventory and analysis of the food plants of southern Africa. South African Journal of Botany 122 (2019) 136-179
- World Checklist of Useful Plant Species 2020. Royal Botanic Gardens, Kew

PROVIDED BY Food Plants International

French, B.R. & Maynard, A.R. Food Plants International Database. Food Plants International

LICENCE CC BY-NC-SA 3.0

ONLINE FACT SHEET <https://www.echocommunity.org/en/resources/ff7f73ce-01f4-405c-bf53-d4c0a692ec47>