

Common names Apocynaceae

# Carandas plum, Simple-spined carissa,

Scientific name

Carissa edulis

**Edible portion** 

Fruit, Roots - spice, Flowers, Leaves,

# **Description**

A shrub or small tree. It grows up to 1.5-5 m tall. It has green straggly branches. There are many spines 10-40 mm long. The spines are simple. The leaves are opposite one another. They have short stalks. The leaves are leathery and widest below the middle. The flowers are white and pink. The are in the shape of a tube and 18 mm long. They are crowded in the axils of leaves. The fruit occur as berries in pairs. They are round and about 1 cm across. They are waxy and dark red when ripe. They have red sweet pulp. The fruit are edible.

Probably now Carissa spinarum

### **Distribution**

It is a tropical plant. It occurs in semi arid lowlands and in the highlands. They occur in regions with temperatures in the range 14-22°C. They occur from sea level to 2400 m but are more common at 800-1200 m altitude. It is damaged by frost but resistant to drought. They will grow on most soils. In Tanzania it grows in areas with a rainfall between 1,000-2,100 mm. They do not do well in wet areas. It can grow in arid places. It often grows on termite mounds. It suits hardiness zones 10-11. In Yunnan.

### Found in:

Africa\*, Angola, Arabia, Asia, Australia, Benin, Botswana, Burkina Faso, Cameroon, Central Africa, China, Côte d'Ivoire, Cuba, East Africa, Eritrea, Ethiopia, Ghana, Guinea, Guinée, India, Indochina, Ivory Coast, Kenya, Madagascar, Malawi, Mali, Mozambique, Namibia, Nigeria, Oman, Pacific, Philippines, Rwanda, Saudi Arabia\*, SE Asia, Senegal, Seychelles, Somalia, South Africa, Southern Africa, South Sudan, Sudan, Tanzania, Thailand, Togo, Uganda, West Africa, Yemen, Zambia, Zimbabwe,

#### Use

The fruit are eaten raw especially by children. They can be made into jam.

The fruit are fermented into wine or vinegar.

The roots are eaten as a boiled vegetable. They are used as a spice.

The leaves are used as a pot herb.

## Cultivation

Plants can be grown from seed or cuttings. For hedges they are planted at between 1 and 2 m apart and pruned to 1-1.5 m high. Seedlings need watering. Seeds are best sown in a nursery and then transplanted. Plants can be used as a hedge and pruned.

### **Production**

Fruit reaches maturity 4 months after flowering. Fruit are collected during the dry season. Fruit are harvested off the tree.



Please Note: Except where otherwise noted, content on this site is licensed under a Creative Commons Attribution 3.0 Licence - this means you can share it freely, as is and with acknowledgement.



# **Nutritional Values**

Carissa edulis

Edible Part	<b>Moisture</b> /100 g edible portion %	Energy KJ	Energy Kcal	Protein g	Provit A μg	Vit C mg	lron mg	Zinc mg
Fruit	87.3	1745	418	9.2				
Root								

FOOD PLANTS INTERNATIONAL	d Plants	Helping the Hungry Feed Themselves Well through the strategic use of God's amazing natural resources