

Potentially Important Food Plants of Bangladesh



**FOOD PLANT
SOLUTIONS
ROTARIAN ACTION GROUP**

*Solutions to Malnutrition
and Food Security*



A Project of the Rotary Club of Devonport North,
District 9830 & Food Plants International

www.foodplantsolutions.org

Potentially Important Food Plants of Bangladesh

Dedication

This book is dedicated to the 3 billion hard working farmers and families around the world who cultivate these, and other, food plants for their own subsistence, and who help conserve them in their rich diversity for other people to enjoy.

Preface

This guide is based on information from the Food Plants International (FPI) database developed by Tasmanian agricultural scientist Bruce French. The source material and guidance for the preparation of the book has been made possible through the support of Food Plants International, the Rotary Clubs of District 9830, particularly the Rotary Club of Devonport North who founded Food Plant Solutions, (previously the Learn ♦ Grow project), and many volunteers who have assisted in various ways.

The selection of plants included in this guide has been developed by Lyndie Kite working in a voluntary capacity using the selection criteria developed by Food Plant Solutions. These selection criteria focus on the local plants from each of the main food groups with the highest levels of nutrients important to human nutrition and alleviation of malnutrition. It is intended as a **Draft Guide only** to indicate some important food plants that serve as examples for this purpose. Other important nutritious plants may be equally useful, and it is recommended that the FPI database be used to source information on the full range of plants known to occur in Bangladesh. This guide has been developed with the best intention to create interest and improve understanding of the important local food plants of Bangladesh, and on the understanding that it will be further edited and augmented by local specialists with appropriate knowledge and understanding of local food plants.

Food Plant Solutions was initiated by the Rotary Club of Devonport North to assist in creating awareness of the edible plant database developed by Food Plants International, and its potential in addressing malnutrition and food security in any country of the world. In June 2007, Food Plant Solutions was established as a project of Rotary District 9830, the Rotary Club of Devonport North and Food Plants International. The primary objective of the project is to increase awareness and understanding of the vast food resource that exists in the form of local plants, well adapted to the prevailing conditions where they naturally occur, and how this resource may be used to address hunger, malnutrition and food security. For more information, visit the website www.foodplantsolutions.org. More detailed or specific information on plants, including references to material by other authors, is available on DVD on request.

Disclaimer: This Field Guide has been produced using information from the “Edible Plants of the World” database compiled by Bruce French of Food Plants International. Although great care has been taken by Food Plants International and Food Plant Solutions, neither organisation, or the people involved in the compilation of the database or this Field Guide:

- makes any expressed or implied representation as to the accuracy of the information contained in the database or the Field Guide, and cannot be held legally responsible or accept liability for any errors or omissions
- can be held responsible for claims arising from the mistaken identity of plants or their inappropriate use
- assume responsibility for sickness, death or other harmful effects resulting from eating or using any plant described in the database or this Field Guide

Always be sure you have the correct plant, and undertake proper preparation methods, by consulting with specialist scientists or local users of the plant. The Food Plants International database, from which the information in this Field Guide is drawn, is a work in progress and is regularly being amended and updated.

Contents

- INTRODUCTION 1**
- STARCHY STAPLES 13**
- LEGUMES..... 23**
- LEAFY GREENS 31**
- FRUIT 39**
- VEGETABLES..... 48**
- NUTS, SEEDS, HERBS AND OTHER FOODS 55**
- NUTRITIONAL VALUES OF FOOD PLANTS BY PLANT FAMILY..... 63**

Introduction

This book is designed as a simple introduction to useful, and sometimes under-utilised, food plants of Bangladesh. It is hoped people will take greater pride and interest in these plants and become confident and informed about how to grow and use them. Many of the local food plants that occur in every country are very good quality foods. Unfortunately, people often reject traditional food plants in favour of introduced varieties. The principle behind Food Plant Solutions is to encourage the use of these local plants.

Local food plants are often very good

People sometimes think that local food plants are not very special and that any food plant that is new or comes from another country must be a lot better. While this is sometimes true, it is often not the case. Many of the newer or introduced food plants, such as the round or ballhead cabbages and lettuce, have very little food value. Many traditional tropical green, leafy vegetables and ferns have 10 times or more food value as ballhead cabbage or lettuce. It is important to find out more information about the food value of different foods if we want to eat well. Citrus fruit, such as lemons and oranges, are often grown for vitamin C that helps keep people healthy. These fruits do not grow well in the tropics - the common guava fruit has three times as much vitamin C and is loved by children. This is just one example showing there are often much better choices of local foods with higher levels of important nutrients.

Growing food

Growing food to feed a family is, without doubt, one of the most important things anyone can do. The more interest you take in your garden and the more you learn about plants and how to grow them well, the more interesting and fun food gardening becomes.

A country with very special plants

The local food plants of most countries have not been promoted and highlighted in the way they deserve. Visiting a local food market will quickly show what a rich variety of food plants can be grown in this country. Good information about these plants is often still in the minds and experience of local farmers, and has not been written down in books. This can make it hard for the next generation of young people to find out how to grow them.

In many countries, some of the traditional food plants are only harvested from the wild and others are only known in small local regions. Others have hundreds of varieties and are the main food for people in different regions. Information on these plants, and in many cases, their food value and the pest and diseases that damage them, is available in the Food Plants International database.

Getting to know plants

People who spend time in gardens and with their food plants get to know them very well. It is a good idea to learn from someone who grows plants well. Each plant grows best in certain conditions and there are often special techniques in getting it to grow well. For example, sweet potato will not form tubers if the soil is too wet, but it may still grow lots of green leaves. Taro will grow in light shade, but sweet potato will not. Ginger can grow in fairly heavy shade. Pruning the tips of betel leaf or pepper vines will cause more side branches to grow and therefore, produce more fruit. Stored yam tubers need special treatment if you want them to put out shoots early. Potatoes should be grown from new seed pieces each year to prevent build up of virus diseases. Tomatoes will yield fewer but larger fruit if lower branches are pruned. There are lots of unique things about every plant and learning about these helps a good gardener produce more food.

Naming of plants

Many food plants have local names, as well as a common English name. Every type of plant also has its own scientific name. Although the scientific name might not be widely recognised, this is the link by which people in different countries and with different languages can recognise the same plant. We know that many plants are grown in many different countries, but relying on local or common names, we might not recognise the same plant grown in different places. By using scientific names to accurately identify plants, we can get useful information from people in other countries. Wherever possible, plants in this book are named by their common English name and their scientific name.

An Important Note

As noted above, a guiding principle behind Food Plant Solutions is to encourage the use of local plants. It is acknowledged that some of these may be major agricultural crops that are already well known, such as:

- Rice
- Oats
- Mango
- Wheat
- Maize/corn
- Banana


There are others, but these serve as examples.

As a general principle, these types of plants are not included in a field guide, as they are well known, and in some cases (e.g. corn), can be relatively high input crops. However, if they are particularly important in a country, such as rice in many Asian countries, they may still be included. The main purpose of the Food Plants Solution project is to look beyond these types of crops, and focus on plants that are less well known and, as noted above, often have superior food value and lower input requirements.

It is also important to note that while some plants have extremely high levels of some nutrients, many of these are used as flavouring foods (condiments), and are generally used only in small amounts. Therefore, the nutritional contribution they make to the diet will be relatively small. Typical of these types of plants are coriander, parsley etc. Once again, these should not be considered as major food sources. A few may be included in the *Nuts, Seeds, Herbs and other foods* section of the field guide.

Nutrient Value

Our bodies need a variety of food plants to enable us to grow, stay healthy and have enough energy to work. Different foods are needed to provide energy, protein, vitamins and minerals. The following diagram highlights the iron content value of some traditional edible, tropical plant leaves, compared with cabbage. Iron is a nutrient that is very important for our bodies and especially our blood. People who are short of iron become anaemic and lack energy.



The relative iron content of some edible leaves

A healthy balanced diet

Good nutrition, or eating a healthy balanced diet, is really very simple. If people eat a wide range of food plants, their bodies will normally get a balanced amount of all the different nutrients they require. If a nutrient is lacking in one food plant, then they are likely to get it from another plant if they are eating a range of food plants. For this reason, everybody should eat a range of different food plants every day. The food group that is especially important for young people is the dark green leaves. Everyone should eat a good serving of dark green leaves every day. They have many vitamins and minerals, as well as protein. There are many spices or flavouring plants that can improve the taste of foods, but taste should be considered separately from food value.

Learning to cook well

Even though some nutrients in food can lose some of their value during cooking, it is normally much safer to cook all food plants, at least for a short time. Bacteria, which cause diarrhoea, can occur in gardens and on food plants. These are killed during cooking. Some plants contain cyanide, a chemical that makes them bitter and poisonous. This happens often with cassava (tapioca, manioc) and beans, but can also occur in many other plants. Boiling the food for two minutes normally destroys cyanide and makes the food safe to eat. Some of the nutrients our bodies need (such as vitamin A for good eyesight) becomes more readily available when food is cooked in oil.

Learning to grow “wild” food plants

Many plants grow wild in the bush and are not cultivated by people. We can normally find someone who has taken an interest in them and has learned to grow them. This may be people from a different region. It may be that in their area they have found better types than the ones that simply grow wild.

Saving better types of plants

If we simply allow plants to grow from seed, the improvements that have been made in finding sweeter or better types may get lost. Some fruit trees are like this and the fruit produced may not be sweet at all. It is often necessary to take cuttings from a tree to be sure the new plant is exactly the same as the old one. If the plants won't easily grow from cuttings simply by sticking a piece of the branch in the ground, (and keeping it watered), there are other ways of helping these plants to form roots and start to grow. One good way is to make a small cut in the bark of a young branch and then wrap soil around the cut and cover it with plastic. With some plants (like guava), new roots will start to grow from this cut and grow into the soil wrapped around the branch. It can then be cut off and planted. This is called air-layering. A similar method is used with the roots of breadfruit. A shallow root is uncovered and a small cut made from which a new sucker will start to grow. This can be cut off and replanted.

Growing from cuttings and suckers

Many food plants are grown from cuttings and suckers. This is very important, as it allows all the different kinds of yams, taros, bananas, sweet potato and sugarcane to be continually grown and ensures the varieties are preserved. Each plant has its own special propagation method. It is important to use healthy planting material, as diseases can be spread in planting material. In many cases, small tubers of yams, taro etc. are stored for planting for the next crop. This is not always good practice, as the small tubers could be the result of diseases (such as viruses) in the plant. A good rule is to take cuttings or save tubers from the best plants for re-planting.

Saving seed

Some food plants are grown from seed. Sometimes this is very easy as the seeds are large, store well, grow easily and grow the same as the original plant. It is more difficult with other plants. Many large fleshy seeds, such as breadfruit, need to be planted while still fresh as they do not store easily. Other seeds do not “breed true” or do not grow into new plants that are the same as the original plants. With many of these plants, it may be necessary to find ways of growing them from cuttings or other methods such as grafting. Some plants “inbreed” and get smaller or poorer. This happens when a plant self-pollinates or receives pollen from a close relative. Corn grown in small plots normally does this and the plants grown from seed grown in this situation get smaller and smaller each year. The seed needs to be saved from several different plants with different history and then mixed together before sowing. All the seeds on one cob are related and will inbreed. Some seeds develop a hard seed coat and need to be scratched, soaked in water, or even put into hot water, before they will start to grow. Saving local seeds is often a good idea as they are already adapted to local conditions. For example, seed saved from pumpkins grown locally will produce plants with less pest and disease damage than those grown from imported seed. This is not to say that imported varieties might not be better, but it is worth checking that the new plant will perform as well as, or better than, those currently grown, before going to the trouble of importing seed. *If you can't get seeds or planting material from local gardens, it is probably not a suitable local plant! Once again saving seed from the better plants helps ensure that you have better crops the next time you plant.*

Growing a garden of mixed plants

In nature, one variety of one plant never grows alone. There are always lots of different plants of different kinds and sizes, all growing together. Anyone who has ever walked into a tropical jungle will know this very well. Growing plants in a food garden in a way similar to how they grow in nature, as a mixed group of plants, is very good agriculture. Mixing plants in a garden usually gives more reliable food production, as any disease from one plant will wash off in the rain onto a different plant, where it cannot survive. Small plants fill the gaps and reduce the need for weeding.

Different types of plants for food security

There is another reason for growing a range of food plants in a local garden or around a village. If something goes wrong, like extreme insect damage to plants, some disease occurring in the garden, or a poor growing season, some plants will be more damaged than others. With a variety of plants, there will still be some food to eat until the other plants recover and grow again. Also, a wide variety of plants will mean that different ones will be maturing at different times, which helps ensure a continuous supply of food. There are shrubs that can be planted as edible hedges around houses, and fruit and nut trees that need to be planted as a gift for your children, several years before they will be able to enjoy them. Some nuts can be stored and eaten when other foods are not available. Most tubers will store well for a few months in the right conditions.

Crop rotation

Crop rotation involves planting different crops in succession to improve soil fertility and reduce the impact of insect pests and diseases. The crops in the rotation should be selected to reduce the risk of carry-over pests and diseases from one crop to the next. A three-year rotation would normally involve growing a legume to increase soil nitrogen levels, and then two different crops before a legume is planted again. Maximum suppression of diseases and insects should be achieved if the non-legume crops are alternated between grass and broad-leafed plant families. Crops planted in a mixture of two or more species will provide greater diversity, and hence reduce losses due to pests and diseases.

Looking after the soil

Gardeners in traditional tropical agriculture usually move their gardens often by shifting to a new piece of land. There are usually three reasons for this:

- In the tropical lowlands, weeds can become a very big problem. There are usually a lot fewer weeds in the first year or two after clearing and burning the land, but weeds increase in the following years.
- Some of the nutrients in the soil are used each year and the soil becomes poorer and plants do not grow as well. There are ways of reducing this loss of nutrients.
- Very small worms called nematodes build up in the soil after a few years and get into the roots, especially of annual vegetable plants, and stop their roots working properly. For example, root knot nematode will cause the roots of plants like tomatoes and beans to become twisted resulting in poor growth of the plant.

Control of soil erosion

Erosion, or loss of soil, occurs when wind or running water carries soil away from cultivated areas. If erosion continues year after year, the land will become unproductive. Trees and shrubs can be planted in strips across the direction of the dominant wind to reduce wind erosion, or across the slope to help slow water that flows over cultivated soils. Growing crops with good leaf cover, or leaving crop residues to cover the soil surface, will significantly reduce soil loss during windy or wet seasons. When possible, any soil disturbance, such as tillage, should occur after the risk of windy weather or wet weather run-off has passed. In steep areas, tillage should be on the contours around hills, rather than up and down the slope. Various techniques can be used to leave ridges or terraces in the tilled soil to help slow water, or reduce the effects of wind, and hence reduce soil loss. Always reduce tillage to the minimum amount possible while still achieving a successful crop.

Building up the soil

When a new garden has been cleared, it has lots of leaf mulch and other old plant material. This provides plant nutrients for new plants to grow. There is a simple rule for growing plants and improving the soil - "If it has lived once, it can live again." Any old plant material can provide nutrients for new plants to grow, but it must be allowed to rot into mulch or compost for this to happen. If this plant material is burnt, some nutrients, especially phosphorus and potassium ("potash"), get left behind in the ashes for new plants to use, although it also allows these important nutrients to be lost by being washed away by rain. But with burning, other important nutrients, such as nitrogen and sulphur, get lost in the smoke and disappear from the garden and soil. These last two plant nutrients are especially important for growing green leaves and when their levels are low, plants grow small or pale green. When nitrogen is lacking, the old leaves of the plant go pale and fall off early, and when sulphur is lacking, the young leaves go pale. Wherever possible, old plant material should be covered with some soil to allow it to rot down and not simply dry out or get burnt. Cutting it into small pieces will help it break down more quickly into usable compost.

Poor soils where crops won't grow

When soils are very acid (or sour), plants cannot get the necessary nutrients. Natural chemicals in the soil that are toxic to plants when present at higher levels become soluble, get into plants, and stop them growing. Adding limestone to these soils can improve them. Using compost will not make them less acid, but will keep the plant nutrients in the soil in a more readily available form that plants can use.

Soil nutrients

Plants need 16 different kinds of plant food or nutrients in different amounts to grow properly. A plant that has already been growing will have these nutrients in them and probably even have them in a balanced amount. That is why composting old plant material is so important. Plants usually show some signs or symptoms if any of these nutrients is running out.

One of the most common and important nutrients for plant growth is nitrogen, which actually comes from the air, but gets into plants through the soil. When plants are short of nitrogen, their older leaves often become yellow or pale. When grass family plants, like corn, are short of nitrogen, the centre of the oldest (lowest) leaves starts to develop a dry or dead V-shape. The plant cannot find enough nitrogen in the soil so it gets it from an old leaf to grow a new leaf. This causes the old leaf to die, forming a characteristic V-shape in the centre of the leaf. The plant does not get any bigger as an old leaf dies each time a new leaf is produced. Village farmers often walk through grassland before they clear it for gardens, looking to see if the grass leaves are dry and dead, because they know gardens on this soil won't grow well. It is necessary to use compost or legumes (such as beans) to put nitrogen back into the soil. Growing plants from the bean family (legumes) is an efficient way to increase the level of nitrogen in the soil.

Corn is a good plant for indicating which nutrients are running short in the soil. If the older leaves go dry along the edges, the soil is running out of potash. If leaves that are normally green develop a bluish colour, the soil is short of phosphorus. Generally, leafy crops need lots of nitrogen, and root crops need lots of potash.

Making compost

Compost is old plant material that has been allowed to rot down into a fine, sweet smelling mulch that is full of nutrients that can be put back on the soil to grow new plants. Compost returns nutrients to the soil, improves the soil's ability to retain moisture and also helps improve soils that are acid or saline.

Making good compost is very simple. A simple heap of plant material can be made in the corner of a garden or near a house. Cutting the plant material (especially stems), into small pieces no longer than about the width of a finger, will help it break down quicker. If possible, make layers of plant material, then a small layer of soil, and then scatter fire ash on top. Keep repeating this process to make a heap. A good compost heap should be warm inside. Be careful with diseased plant material. This should be burnt, otherwise the disease may be spread when you use the compost at a later date.

The composting process is carried out by small bacteria that live in the soil and feed on decaying plants. They break down old plant material into compost. These bacteria are living, so they need air, water and food. A good compost heap must have air, so don't cover it with plastic or put it in a container. This makes a foul smelling compost, as different bacteria that don't need air turn it into an acid mixture that preserves it. Good compost must have moisture, so keep the heap damp, but not too wet. The compost bacteria like a balanced diet, which means that both green material and dried material is needed to balance the carbon and nitrogen in the compost pile. If the compost material gets too dry and brown, it will not break down, and if it gets too green, it will go slimy. Using a little bit of compost from an old heap will make sure the right bacteria are there to start the whole process off.

As soon as the plant material is broken down to a fine mulch it can be put onto the garden. It is best if it is dug in, but if it is regularly put onto the surface of the garden, worms will mix it into the soil. A handful of compost placed at the base of each plant can be beneficial.



Pests



There are a large number of insects that enjoy sharing our food with us! We should not try to kill all these insects as they have an important role to play in keeping everything in nature in balance. What we need to do is to learn to manage these insects so we can all get some food to eat! Some insects are attracted to lights, and if the garden is near village lights some insects can cause a lot of damage. If large areas of one particular crop are planted, insects can breed more quickly and cause a lot of damage. As an example, insects called armyworms can breed up in large numbers on the shade trees of cacao and then move “like an army” into gardens. Some insects are large and breed slowly and can be picked off and removed. The large, green grubs with pointy tips that hide under taro leaves are best controlled by simply picking them off. Some insects, like taro beetles, can be a serious problem, but the young curl grubs of this insect are tasty if you catch and cook them. Some insects do not like sunlight. The very small moth that damages banana fruit is like this. Simply pulling off the leafy bracts over the banana fruit reduces the damage, as this lets sunlight in and the insect flies away. The best rule for reducing pest damage is to grow healthy plants, as they suffer less damage. Spiders, ladybirds, hover flies and many other insects also feed off the insect pests that attack our crops and should be encouraged.


Diseases

The living organisms that cause disease are much smaller than insects. These disease organisms can often only be seen with a microscope. There are three main kinds of disease organisms - fungi, bacteria and viruses. Fungi are like the mushrooms we eat, only very much smaller. They usually make distinct dry spots on leaves and other plant parts. Fungi have spores that often blow in the wind. Bacteria are often smaller and live in damp places. They usually make plants go soft and squashy, and they may cause a smell. Bacteria are mostly spread with rain and in water. Viruses are very, very small and usually make irregular stripes and patterns on leaves and other plant parts. Viruses usually spread in planting material or in the mouths of small sucking insects. Plants infected by viruses are often yellow, and may be stunted, or have curled or unusual shaped leaves. One common fungus disease on sweet potato causes the leaves to become wrinkled and twisted. It usually gets worse in old gardens and where soils are running out of nutrients. It doesn't affect all kinds of sweet potato to the same extent. The answer is not to stop the disease, but to improve the soil. The general rule is that healthy plants that are growing well will suffer less damage from disease.

Food value charts for a selection of plants from Bangladesh







Note regarding plant selection: In compiling these field guides, we acknowledge that some staple foods and commercial crops which are grown widely in the target country may be omitted. Such foods are often in the starchy staple category (e.g. rice, corn). This does not mean that they are not useful, but merely reflects a desire for the Food Plant Solutions project to concentrate on plants that are less well known and/or underutilised.

Starchy staples

English: Comb fringe grass

Scientific name: *Dactyloctenium aegyptium*

Local:

Plant family: POACEAE

Description: An annual grass. The stems are slender. They can lie along the ground. These can form roots at the nodes. They can have runners and form mats. It is 15-60 cm high. The edges of the leaf sheaths have small hairs. The leaf blades are flat and 5-20 cm long by 0.2-0.6 cm wide. The surfaces are lumpy/hairy. It tapers to the tip. The flowers spread like fingers on a hand. There are 2-9 flower stalks. They are long and narrow. They often spread out horizontally. The spikes are on one side of the stalk. The tip is bare. The seed grains are about 1 mm across.



Distribution: A tropical plant. It grows in disturbed weedy places especially on sandy soils in S China. It grows in tropical to warm temperate regions. It grows on clayey, sandy or black soil along the borders of ponds, swamps and bogs. In West Africa it grows from sea level up to 2,000 m altitude. It grows in alkaline and salty soils. It grows in areas with an annual rainfall between 100-1,580 mm. It can grow in arid places.

Use: The seeds are husked then boiled into a porridge. They are also roasted in a hot pot to soften them. It is then pounded into flour and cooked into porridge. The rhizome or runners are eaten raw.

Cultivation: Plants can be grown from seeds.

Production: The seeds are collected during the dry season. The seeds can be stored for several months.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	7.5	1234	9.8	-	-	6.9	4.7

Image sourced from: Giles Tran AFZ @feedipedia.org

Starchy staples

English: Finger millet

Local:

Scientific name: *Eleusine coracana*

Plant family: POACEAE

Description: An annual millet grass. It is robust and forms many tillers or young shoots from the base. It grows 40 - 120 cm tall. The stems are somewhat flattened and the leaves are narrow. The flower heads are made up of 2 - 7 finger like spikes, 1.5 cm across and 10 - 15 cm long. These in turn have about 70 smaller spikes. Each one of these smaller spikes has 4 - 7 seeds. The seeds are roughly rounded and 1 - 2 mm across. There are *coracana* and *africana* subsp.



Distribution: It is a very drought resistant tropical plant.

For good yields, it needs good soil drainage and adequate moisture. It cannot stand water-logging. It is an important crop in areas where annual rainfall is 900 – 1,250 mm. It especially suits areas with long hot summers. It needs a minimum temperature above 18°C and does best where temperatures are above 27°C. It grows from sea level to 2,400 m altitude in Africa. It is a short day length plant and does best where daylength is 12 hours. It can grow in arid places.

Use: The seed are eaten either roasted or ground into flour. This is used for porridge and flat bread. Alcohol is brewed from the grain. The leaves are also edible.

Cultivation: It is grown from seed. Often plants are grown mixed with sorghum or maize. Good soil preparation is needed to reduce weed competition. Seed can be broadcast or drilled. Young plants need to be weeded and thinned. Seed viability drops to about 50 % after 2 years. Spacings of 5 cm apart in rows 30 - 33 cm apart, or 10 - 12 cm apart in rows 25 cm apart are recommended. About 25 - 35 kg of seed per hectare are needed if seed are broadcast. 5 - 10 kg per hectare are required if seed are drilled. Using fertiliser can dramatically increase yield. 125 kg per hectare of sulphate of ammonia when plants are 15 cm high is used in Uganda.

Production: It is self pollinating and pollination occurs over 8 - 10 days. Millet seed stores very well and can be stored without damage for 10 years. Often it is stored on the head. Yields of about 450 - 900 kg of dried grain per hectare are usual. This can easily be increased to 1,650 kg per hectare. Crops take 3 - 6 months until harvest.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seeds	11.7	1594	6.2	-	-	5.3	-

Starchy staples

English: Barley, Pearl barley

Scientific name: *Hordeum vulgare*

Local:

Plant family: POACEAE

Description: An erect annual grass. It grows 80-120 cm tall. The nodes are solid and the internodes are hollow. The leaves are narrow. There are 5-10 leaves. They are produced alternately on opposite sides of the stem at the nodes. The leaves are narrowly sword shaped and 5-40 cm long by 0.5-1.5 cm wide. The flowers are greenish. Flowers have long awns. The fruit is a grain. It is oval and narrow. There are a range of named cultivated varieties.



Distribution: A temperate plant. It requires full sun and well-drained soil. It can tolerate saline conditions. In Nepal it grows to 3,500 m altitude. It grows in the mountains in Java.

Use: The grains are eaten. They are mixed with wheat for chappatis. They are also used in soups. They are also used for bread and breakfast cereals. They have a low gluten content. The sprouted seeds are eaten in salads. Barley water is made by soaking the barley in water and flavouring with lemon. The young seedlings are dried and powdered and marketed as *barleygreen*. Roasted seeds are added to coffee. The seeds are also soaked until they sprout and produces malt. This is used for alcohol production.

Cultivation: Plants are grown from seed. Seed can be planted 2-6 cm deep. Often 200-250 plants are grown per square metre.

Production: It has a relatively short growing season and matures quickly.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed(boiled)	69.6	510	2.7	-	-	6.0	-
seed	13.7	1367	10.5	-	-	6.0	-

Starchy staples

English: Cassava

Local:

Scientific name: *Manihot esculenta*

Plant family: EUPHORBIACEAE

Description: A plant which can re-grow year after year from the thickened roots. It has several stems. The stems are woody and have some branches. Plants grow up to 3 metres tall. Stalks have distinct scars where leaves have fallen. The leaves tend to be near the ends of branches. The leaves are divided like the fingers on a hand. The leaves have long leaf stalks. The leaves have 3 - 7 long lobes which can be 20 cm long. These are widest about 1/3 of the distance from the tip and taper towards the base. The colour varies. It produces several long tubers. These can be 50 cm long by 10 cm across. The flowers are on short stalks around a central stalk. They are produced near the ends of branches. The female flowers are near the base of the flower stalk and the male flowers higher up.



Distribution: A tropical plant. Plants grow from sea level up to about 1,650 m. In Fiji they grow to 900 m. They can grow in poor soil and can survive drought. It is native to tropical America. It grows between 25°N and 25°S and needs a rainfall above 750 mm. It suits hardiness zones 10 - 12.

Use: The tubers are eaten after thorough cooking. They are boiled, roasted or made into flour. The starch is used in puddings, soups and dumplings. Young leaves are edible after cooking. They are also sometimes dried and stored. Seeds are also eaten.

Caution: Bitter kinds of cassava contain poison but this is destroyed on heating. This kind of cassava should be cooked, sun dried, soaked and cooked again.

Cultivation: Cassava is planted from sections of the stalk. Sections about 15 - 20 cm long of the more mature woody stem are cut and stuck into the ground. They can be completely buried or put at almost any angle and it affects the growth little. Soon roots form and leaves start to sprout from the stalk. Cassava seeds need a soil temperature of 30°C for their germination. Flower and fruit production is more common under lower temperatures such as in highland or less equatorial conditions.

It is not necessary to dig a hole to plant cassava and on many soils where the soil is loose it can be planted without digging the soil first. Cassava does not suit waterlogged soils and preferably they should not be too shallow or stony.

Cassava can be planted at any time of the year but to get started it needs moisture so is often planted near the beginning of the wet season. The crop once established can survive for several months without rain. The ability to tolerate drought varies significantly with cultivar. During drought less and smaller leaves are produced and leaves die off more quickly but storage roots can be increased in the short term.

Because cassava can still grow satisfactorily in poorer soils it is often put last in a rotation after others crops have already been grown on the piece of land. Cassava is more responsive to nitrogen and potassium than phosphorus under many field situations. Nitrogen can increase cyanide levels. Under very acid conditions with high soluble aluminium levels, cassava has been able to achieve and maintain top growth but with significantly reduce root yields. When drainage is good and soil moisture is adequate, cassava stalks can be planted at any orientation from horizontal to vertical, but in very sandy soils horizontal planting is best and in heavy clay soils vertical planting is best.

Because of the slow growth in early establishment stages, soil loss from erosion with heavy rains can be significant. To avoid this planting should be timed so that the maximum vegetative growth is occurring during the heaviest rains. A leaf area index between 2.5 - 3.5 is optimal for cassava yield. The critical period for weed control is the time from 2 - 8 weeks after planting. Cassava tuber bulking is delayed under shaded conditions. Yields are also reduced. In mixed cropping situations using crops which mature early, allowing the cassava time to recover, is one possible strategy. For optimum production shading should be avoided.

Cassava takes about 10 - 12 months to produce mature tubers in the lowlands tropics although some varieties produce a smaller yield earlier. Yields in the range of 20 - 45 t/ha have been recorded for 12 - 14 month crops. The plants can be left growing and the tubers stored in the soil for considerable time. Crops of 24 months duration occur. Once the tubers have been dug they do not keep for more than a few days. Pre-harvest pruning of plants increases the storage time of tubers after harvest.

Spacing and plant density varies with soil climatic conditions and variety. Plant densities from 10,000 to 30,000 plants per hectare are used. Plants from the higher density crops have been shown to have quick post harvest deterioration. Mulching has given significant yield increases in some conditions. It also reduces the incidence and damage of some root boring insects.

Production: Plants can be harvested after 10 months in the lowlands. There are some faster growing varieties. Yields in the range of 20 - 45 t/ha have been recorded for 12 - 14 month crops.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	62.8	625	1.4	30	15	0.23	0.48
leaf	82.0	382	7.1	57	275	7.6	-

Starchy staples

English: Taro

Local:

Scientific name: *Colocasia esculenta*

Plant family: ARACEAE

Description: This plant has large flat leaves on the end of upright leaf stalks. It grows up to 1 m high. The leaf stalk or petiole joins the leaf towards the centre of the leaf. The leaves are 20 - 50 cm long. Near the ground a thickened rounded corm is produced. Around this plant there is normally a ring of small plants called suckers. Many different varieties occur. If left to maturity, a lily type flower is produced in the centre of the plant. It has a spathe 15 - 30 cm long which is rolled inwards. The flowers are yellow and fused along the stalk. There are many named cultivated varieties. Taro comes in two basic forms. The Dasheen type *Colocasia esculenta* var. *esculenta* and *Colocasia esculenta* var. *antiquorum* or the Eddoe type.



The basic difference is the adaptation of the Eddoe type to storage and survival in seasonally dry places, while the dasheen type needs to be maintained in a more or less continuously growing vegetative stage.

Distribution: It is a tropical plant. Taro grows from sea level up to about 2,300 m altitude in the tropics. It grows well in humid places. It can stand damp soil and grow under light shade. It suits hardiness zones 9 - 12.

Use: The corms, petioles and leaves are all edible after cooking. The leaves are also dried and stored. Fresh leaves can be stored for 4 - 5 days.

Caution: Some varieties burn the throat due to oxalate crystals.

Cultivation: Taro can be planted from cormels or from the top of the central corm. Other sections of the corm could also be used but this is not commonly done. Flowering of taro and seed production can lead to new cultivars. Flowering can be promoted by the use of gibberellic acid. The general growth pattern is for an increase in top growth, in terms of leaf number, leaf area and petiole length, to continue for about 6 months under tropical lowland conditions then for each of these to decrease and tuber storage to continue to increase. Corm weight increases significantly from 5 - 11 months. Starch content also increases with time but protein content declines over the corm development period.

Taro can be grown under flooded conditions but root rots develop if the water becomes stagnant. For flooded cultivation, the land is cleared, ploughed, cultivated and puddled. The aim is to get a field that is flat with embankments allowing the impounding of water. Planting is done into 2 - 5 cm of standing water.

For dryland taro, the soil is prepared by digging, unless a fresh bush fallow is used where the natural friability of the soil allows plants to be put into the undug soil in a small hole that is prepared. Plants are put into a hole 5 - 7 cm deep or deeper. Mulching to conserve moisture and reduce weed growth is beneficial. Setts from corms normally give higher yield than that from cormels. The greater leaf area and root production may be responsible for this. Setts of about 150 g are optimum.

The time of planting is primarily determined by the availability of moisture. Planting is done shortly after the rainfall has become regular, if seasonally distinct wet and dry occur. Higher rainfall, higher temperatures, and higher hours of sunlight, enhance production and determine seasonality of production.

Evapotranspiration for flooded taro averages about 4 mm per day, ranging from 1.5 - 7.2 mm, with a total of about 1,200 mm for the crop. Intermittent moisture can result in irregular shaped corms. Flooding has been found to be more effective than sprinkler irrigation, or furrow irrigation. Increased suckering, giving greater leaf area, seems to be the reason for this.

Taro is sensitive to weed competition throughout most of its growth, but it is more critical during early growth up to 3 - 4 months. About 7 - 9 weedings are required, to keep the crop clean under tropical lowland conditions, where flooding is not used. Due to the decrease in height and leaf area towards the end of the growth cycle when starch accumulation in the corms is maximum, weed competition and weed control are again significant. Mechanical weeding needs to be shallow to avoid damaging the superficial taro roots. A range of herbicides have been recommended in various situations.

Taro produces the highest dry matter yield under full sunlight, but it can still grow under moderate shade. Under shaded conditions it grows more slowly and develops fewer cormels. They require good moisture conditions and have little tolerance for drought. Taro residue has an allelopathic factor which can reduce the germination and growth of other plants, for example, beans.

Taro tends to demand high fertility, and is responsive to additional NPK fertiliser. Higher doses of K increases starch content and higher doses of N increases protein content. Both N and K applications increase oxalic acid content of the tubers.

Spacing affects total yield, and marketable, harvestable yield, of corms. Close spacing increases the corm yield per area, and the shoot yield per area, but decreases the corm yield per plant, and the contribution of sucker corms, to the yield. Where spacings of 30 cm x 30 cm are used, giving about 110,000 plants per hectare, a very large amount of planting material is required, which reduces the net return per unit of planting material. A spacing of 60 cm x 60 cm is more common. Wider spacings of 90 cm x 90 cm reduces overall yield.

Production: Crops mature in 6 - 18 months. Yields of 5 - 15 tonnes per hectare are probably average.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
root	66.8	1231	1.96	3	5	0.68	3.2
leaf	85.0	210	5.0	57	90	0.62	0.7
leaf stalk	93.0	101	0.5	180	13	0.9	-
leaf (cooked)	92.2	92.2	2.7	424	35.5	1.2	0.2

Starchy staples

English: Niger seed

Local:

Scientific name: *Guizotia abyssinica*

Plant family: ASTERACEAE

Description: An erect branched herb. It grows 30-180 cm tall. The stems are soft and hairy. The leaves are usually carried opposite one another. The leaves do not have stalks and they clasp the stem. The leaves have teeth along the edge and the surface is a little rough. The flower head is made up of many small flowers each capable of producing a seed. The fruit (called seeds) are black angled structures. They are up to 12 mm long. The seed inside is 3.5-5 mm long. There are about 250-300 seeds per gram.



Distribution: It is grown in both temperate and tropical zones. It can compete well with weeds. It has some salt tolerance. The temperatures are between 16°C and 20°C where Niger does best. It cannot tolerate temperatures above 28°C average and must have temperatures above 6°C. In Africa it grows between 300 and 2,300 m altitude but does best at 1800-2000 m altitude in Ethiopia. The rainfall is 100-1,300 mm per year where it grows most. At lower altitudes a lower rainfall is satisfactory if it is spread through the growing season. It grows on a wide range of soils. It can grow on poorly drained soils. It needs short day lengths for flowering. It grows in open places. It can grow in arid places.

Use: The seeds can be fried and used as snacks. They are also used in sauces. The seed cake is mixed with honey to make a sweet bread. The seed oil is edible. It is used in cooking.

Cultivation: It is grown from seed. It is often sown mixed with finger millet. Seeds can be broadcast or planted in rows. 10-15 kg of seed per hectare are used for broadcast crops. When planted in rows 5-8 kg per hectare of seed are used. Fertilisers do not improve seed yield much. As petals drop, seeds are harvested to avoid seed loss. Stems are cut near the ground, then dried for a few days before threshing.

Production: Flowering occurs 3 months after sowing. Seeds are ready for harvest 4-6 weeks later. It can take 5 months at altitudes over 2,000 m. Yields can be 300-700 kg per hectare. Yields of 1,400 kg are possible.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	6.2	2019	17.3	-	-	-	-

Image sourced from: thebloomingauction.com

Starchy staples

English: Sweet potato

Local:

Scientific name: *Ipomoea batatas*

Plant family: CONVOLVULACEAE

Description: This is a root crop which produces long creeping vines. The leaves are carried singly along the vine. Leaves can vary considerably from divided like fingers on a hand to being entire and rounded or heart shaped. Purple trumpet shaped flowers grow at the end of the vine. Under the ground fattened tubers are produced. There are a large number of varieties which vary in leaf shape and colour, tuber shape, colour, texture and in several other ways.



Distribution: A tropical and subtropical plant. They grow from sea level up to about 2,700 m altitude in the tropics. Plants can grow with a wide range of rainfall patterns and in different soils. Plants are killed by frost and can't stand water-logging. Plants grow well with temperatures between 21 - 26°C. It can grow with a pH between 5.2 - 6.8. Sweet potato are not tolerant to shading. Under shaded conditions, both foliage growth and storage root production are decreased. Some cultivated varieties can be selected for increased production under mild shade but not heavy shade. The survival of cuttings at planting is also reduced under shaded conditions. Under shaded conditions plant become more climbing and with fewer leaves which are however larger. With increasing shade less tubers are produced and these grow more slowly. Sweet potato tends to be responsive to potassium fertiliser. cultivated varieties are often selected for yield under low fertility conditions. Under lowland conditions in the tropics sweet potato tubers undergo active tuber enlargement from 6 - 16 weeks. Weed control is essential especially during early stages of growth. The rate of ground coverage by foliage varies greatly with growing conditions and cultivar but once ground coverage has occurred weed control is less of a problem. Sweet potato tuber initiation is subject to aeration in the soil. Either heavy clay soils, waterlogged conditions or other factors reducing aeration can result in poor tuber production. For this reason sweet potatoes are often grown on mounded beds. It suits hardiness zones 9 - 12.

Use: Tubers are boiled or baked. They can be steamed, fried, mashed or dried. They can be fermented into alcoholic drinks. They can also be used in pies, cakes, puddings and candies and jams. They can be used in noodles. The chopped and dried tubers can be boiled with rice or ground into flour and mixed with wheat flour to make cakes or bread. The young leaves are edible.

Cultivation: Vine cuttings are used for planting. In grassland soils it is grown in mounds, ridges or other raised beds. In bush fallow, it is mostly planted in undug loose soils. It needs a sunny position. Tubers won't form if the ground is waterlogged when tubers start to develop. Sweet potato is grown by cuttings of the vine. About 33,000 cuttings are required per hectare. These weigh about 500 kg. Vine lengths of about 30 cm are optimum. As long as the vine is adequately inserted in the soil, the length of vine inserted does not significantly affect yield. Fresh sweet potato seeds germinate relatively easily and lead to continuous production of new cultivars under tropical conditions. Excess nitrogen restricts storage root initiation and therefore excess leaves are produced without significant tuber yield. Dry matter percentage increases with increasing age of the crop. Higher dry matter tubers are normally preferred.

Sweet potato are not tolerant to shading. Under shaded conditions, both foliage growth and storage root production are decreased. Some cultivars can be selected for increased production under mild shade but not heavy shade. The survival of cuttings at planting is also reduced under shaded conditions. Under shaded conditions plant become more climbing and with fewer leaves which are however larger. With increasing shade less tubers are produced and these grow more slowly. Sweet

potato tends to be responsive to potassium fertiliser. Cultivars are often selected for yield under low fertility conditions.

Under lowland conditions in the tropics sweet potato tubers undergo active tuber enlargement from 6 - 16 weeks. Weed control is essential especially during early stages of growth. The rate of ground coverage by foliage varies greatly with growing conditions and cultivar but once ground coverage has occurred weed control is less of a problem. Sweet potato tuber initiation is subject to aeration in the soil. Either heavy clay soils, waterlogged conditions or other factors reducing aeration can result in poor tuber production. For this reason sweet potatoes are often grown on mounded beds. In well drained or high organic matter soils digging or mounding is not as essential. Leaf scab (*Elsinoe batatas*) can significantly reduce yield especially in sites where leaf production is low due to low soil fertility. To reduce sweet potato weevil damage plants need to be hilled or have the tubers well covered with soil. Cracking soils can allow the weevil access to tubers.

Production: The time to maturity ranges from 5 months to 12 months depending on the variety planted and the altitude at which it is being grown. Yields range from 6 - 23 t/ha.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber (baked)	72.9	431	1.7	961	24.6	0.5	0.3
tuber (raw)	70.0	387	1.2	709	25	0.7	0.4
tuber (boiled)	72.0	363	1.1	787	15	0.6	0.3
leaf	86.3	168	3.9	105	58	2.9	-

Legumes

English: Bottle gourd

Local:

Scientific name: *Lagenaria siceraria*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is an annual vine with large leaves. It can grow 3 - 9 m long and spread 3 - 6 m wide. The thick stems have furrows along them. It can climb over logs by attaching the tendrils which grow out of the stem near the leaf. The leaves are large and have soft hairs especially underneath. Flowers of both sexes are borne in the same plant. The plant produces male flowers first and these are on long stalks. Next it produces female flowers on short stalks. Flowers are large and white. They can be 10 cm across. They are mainly pollinated by insects. Fruit vary in shape and can be 8 - 90 cm long. They have brown seeds in a whitish green pulp. There are several varieties.



Distribution: A tropical plant that grows from sea level up to 2700 m altitude in the tropics. It grows best in a warm humid climate. It is sensitive to frost and prefers full sunlight. It grows best with a night temperature of 17 - 23°C and day temperatures of 28 - 36°C.

Use: The young fruit are boiled as a vegetable. The skin and seeds are removed and can also be steamed, fried or pickled. Young tips and leaves are edible. They are often cooked with milk or coconut milk to improve the flavour. They are also mixed with other edible leaves. The seeds are sometimes eaten and provide an edible oil. Old fruit are used as containers, and the seeds are not normally edible.

Cultivation: To achieve fast and uniform emergence, seed should be soaked overnight. Seeds are best sown in raised beds. Seedlings emerge in 5 - 7 days. Seedlings can be transplanted is required. Because plants cross pollinate, plant and fruit types vary. Removing the young fruit to use as a vegetable will prolong the life of the plant. Large fruit can be obtained by removing some of the small fruit. A spacing of 1 - 2 m is suitable. It prefers a trellis to climb. Because it is shallow rooted, weeding needs to be done carefully.

Production: It is fast growing and flowers 2 months after seeding.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
bean (dry)	3.2	2399	28.2	-	-	5.3	-
leaf	83.0	180	4.4	66	-	7.4	-
fruit	93.0	88	0.5	25	10	2.4	-

Legumes

English: Pigeon pea

Local:

Scientific name: *Cajanus cajan*

Plant family: FABACEAE

Description: An upright perennial shrubby legume that can live for 3 - 4 years. They can grow up to 4 m tall and spread to 1.5 m wide. It has a bushy appearance and a strong deep taproot. The root nodules are round and sometimes lobed. The leaf consists of 3 narrow, green leaflets which are silvery-green underneath. The end leaflet is larger with a longer leaf stalk. The pea shaped flowers are red and yellow and occur on branched flower stalks which stick upwards in the axils of leaves. Pods are long, straight and narrow, often with 4 - 8 seeds. Seeds vary in shape, size and colour. The pods are slightly hairy. Pods are often 4 - 8 cm long and have a beak at the end. Pods are constricted between the seeds. Many varieties of pigeon pea occur. Some are dwarf and day length neutral.



Distribution: A tropical plant that requires a tropical or subtropical climate. Plants grow from sea level up to about 1,800 m in the tropics. They can tolerate drought and are suited to a drier climate. They can grow in places with less than 600 mm rainfall per year. They do less well in the wet tropics. They suffer in waterlogged soils and are damaged by frost. It can also tolerate heat. It will grow on poor soils cannot grow on salty soils. It can grow in arid places and suits hardiness zones 10 - 12.

Use: Young leaves, shoots and pods are eaten. The pods can be used in curries. The leaves and shoots as potherbs. Young seeds are cooked and eaten like peas. Ripe seeds are also cooked and eaten in soups and curries. Bean sprouts can be produced and eaten. Preparation of the seeds for dahl is somewhat complicated.

Cultivation: They are grown from seeds. It is best to sow seeds where the plants are to grow. Seeds normally germinate easily and well. Before sowing seed it helps to soak them in cold water for one day. Seeds store well if kept cool and dry. A spacing of 1.5 m x 1.5 m is suitable. Plants can be cut back and allowed to re-grow. Plants can also be grown from cuttings.

Production: Plants are fast growing. Pods are ready after 5 months. Mature seeds take about 8 months. Plants will often live for 3 - 4 years. Plants are cross pollinated by insects, or self pollinated.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	10.0	1449	19.5	55	-	15.0	-
pod (young)	64.4	477	8.7	-	-	2.0	-
seed (young, boiled)	71.8	464	6.0	2	28.1	1.6	0.8

Legumes


English: Jack bean

Scientific name: *Canavalia ensiformis*

Local:

Plant family: FABACEAE

Description: A perennial climber, although short kinds do occur. Often it is a more bushy plant than the sword bean. Plants grow up to 1.5 m long. Stems can be hairy. Leaves have 3 leaflets. The leaflets are oval and 5.7 - 20 cm long by 3.2 - 11.5 cm wide. The leaf tends to be wedge shaped at the base. The leaf stalks are 2.5 - 11 cm long. Flowers are red/purple. They occur on flower clusters 5 - 12 cm long and with flower cluster stalks which are 10 - 34 cm long. The individual flower stalks are 2 - 5 mm long. Pods are long and sword shaped. Pods can be 15 - 35 cm long. Seeds are white with a light brown hilum half as long as the seed. Seeds are 2 cm long, by 1 cm across.



Distribution: It grows in tropical and subtropical places. It requires a fairly high temperature (15° - 30°C). It will possibly grow up to 900 m altitude. It is fairly drought resistant and also has some resistance to water-logging and salt in the soil. It can tolerate shade. It can tolerate pH from 4.5 - 8.0 but does best at about 6.1. The optimum mean annual temperature is 14.4° - 27.8°C. Seed germinate between 24 - 27.5°C. It is a short day plant growing well with a daylength of 10 - 12 hours of sunlight. It can grow in arid places.

Use: The leaves and top shoots are eaten. The very young pods are boiled and eaten. The flowers can be eaten. The young seeds are eaten boiled, roasted, or peeled and cooked. The seeds are also fermented. The ripe seeds are roasted and used as a coffee substitute.

Caution: The ripe seeds can contain poison and need to be well cooked and the water changed before eating. They are also often left under running water or fermented.

Cultivation: It is grown from seeds. Seeds need to be 2 cm deep. A spacing of about 60 cm is suitable. Plants preferably need a support to climb over. It benefits from a fertile soil but adding nitrogen depresses yield.

Production: Green pods are produced in 3 - 4 months, but ripe seeds need 6 - 9 months. Yield of seeds can range from 700 - 5,400 kg/ha.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	10.0	1423	20.4	160	0	4.9	-
pod (fresh)	88.0	155	2.4	-	-	-	-

Legumes

English: Lablab bean

Local:

Scientific name: *Lablab purpureus*

Plant family: FABACEAE

Description: A climbing bean which can have vines 1 - 5 m long. It keeps growing from year to year. The stems can be smooth or hairy. Leaves are made up of 3 almost triangular leaflets. The leaflets are 5 - 15 cm long and 3 - 14 cm wide. The side leaflets are somewhat asymmetrical. Often the plants are flushed purple. The flowering clusters are 5 - 20 cm long. Flowers are often white but can vary from red to blue. The pods are flattened, pointed and up to 12 cm long and 2 cm wide. They can be green, purple or white. Inside there are 3 - 5 white or dark seeds. Seed pods have a wavy margin.



The seeds are 0.5 - 1.5 cm long. (This bean is similar to Lima bean but the keel of the flower is not spirally twisted, the pod ends more bluntly with a long thin style at the end and the hilum on the seed is longer.)

Distribution: It is a tropical and subtropical plant. It mostly grows between 750 and 2175 m altitude in the tropics. It is drought resistant and can grow in quite low rainfall areas. Some varieties are short day and some are long day kinds. It suits hardiness zones 9 - 12.

Use: The young pods, ripe seeds and young leaves are edible, cooked. Flowers can be eaten raw, steamed or added to soups and stews. Dried seeds can be cooked as a vegetable. The seeds can also be sprouted then crushed and cooked. The large starchy root is edible.

Caution: Many types can be poisonous. They should be boiled and the cooking water thrown away.

Cultivation: Seeds are sown at 30 x 60 cm spacing near stakes or trees. About 20 kg of seed per hectare are required. Fertilising with nitrogen and potash until flowering is recommended.

Production: Young pods are ready 4 - 6 months after planting and seeds 6 - 8 months. Pods are often harvested over 2 or 3 years. Pollination and seed setting are reduced in cold weather.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	10.0	1428	22.8	-	-	9.0	-
seed (young)	86.9	209	3.0	14	5.1	0.8	0.4
pod (fresh)	86.7	203	3.9	-	1	2.4	-

Legumes


English: Horse gram

Scientific name: *Macrotyloma uniflora*

Local:

Plant family: FABACEAE

Description: A climbing or twining herb. The young growth has white hairs. The stems are hairy. The leaves have 3 leaflets. The leaves are 3.5 cm long. The leaflets are 1-2.5 cm long by 0.7-3 cm wide. They are broadly oval. They are thin textured and with a rounded tip. The flowers are about 0.8 cm across. They are greenish-yellow. There are 1-5 flowers in a cluster in the axils of leaves. The fruit is a pod 3-5.5 cm long by 0.4-0.8 cm wide. They are slightly hairy and dark brown when ripe.



Distribution: It is a tropical plant. It grows in northern Australia. In tropical Queensland it grows from 500-760 m altitude. It needs freely drained soil. It needs a temperature of 20-30°C. It cannot tolerate frost. It is drought resistant and can grow in areas with a rainfall of 900 mm per year. It needs a pH 5-7.5. It can grow in poor soils.

Use: The seeds are parched then boiled, fried or used in curries and soups. The seeds also yield an edible oil. The flour can be processed into sweetmeats. The fleshy root can be roasted and eaten. The pods are cooked and eaten.

Cultivation: Plants can be grown by seed or cuttings. The seeds are sown 1-2.5 cm deep.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	9.7	1392	-	22.5	-	-	-

Image accessed from: Ian Staples, QLD DPI&F Tropical Forages

Legumes

English: Mung bean

Local:

Scientific name: *Vigna radiata*

Plant family: FABACEAE

Description: An upright hairy bean plant which can grow to 1 m tall. It has many branches. The leaves have 3 leaflets, are dark green and grow on long leaf stalks. There are oval stipules at the base of the leaf. Flowers are pale yellow and small. They occur in bunches of 10 - 20 on the ends of long hairy flower stalks. Pods are black and straight. They do not have a beak. Pods contain 10 - 20 seeds which are usually green or golden yellow. They are smaller than black gram. The beans can be black. They have a flat white hilum. There are 2,000 varieties.



Distribution: A tropical and subtropical plant. The plant will grow from sea level up to about 2000 m in the tropics. It is drought resistant but can't stand water-logging. Plants are damaged by frost. They cannot stand salinity. Rainfall at flowering is detrimental. It requires a deep soil. Both short day and long day varieties occur. It can grow where annual temperatures are from 8 - 28°C. It can tolerate a pH from 4.3 - 8.1. It suits a drier climate and can grow in arid places. It suits hardiness zones 10 - 11.

Use: Seeds are eaten ripe, raw or roasted. They are added to soups and stews. They are also fermented. Young pods and leaves can be eaten. The seeds can be germinated for sprouts and used in salads and stir-fried dishes. The seeds are ground and used for starch to make noodles.

Cultivation: Plants are grown from seed. In some areas these are broadcast while for small plots often 2 - 3 seeds are sown in holes 50 - 60 cm apart. Seeding rates of 6 - 22 kg per ha are used in different locations. It normally requires phosphorus fertiliser for adequate growth. Seeds germinate in 3 - 5 days.

Production: Green pods are ready after about 2 months and ripe pods may take another 1 - 2 months. For ripe beans the whole plant is harvested and dried before threshing. Yields of 450 - 560 kg/ha of seeds are common.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	11.0	1432	22.9	55	4	7.1	-
seed (cooked)	-	439	7.0	2.4	1.0	1.4	-
seed (sprouted)	90.4	126	3.0	2	13.2	0.9	0.4

Legumes

English: Chickling Vetch

Local:

Scientific name: *Lathyrus sativus*

Plant family: FABACEAE

Description: An annual herb. It is a scrambling plant. It grows 100 cm high and spreads 100 cm wide. The stems are winged. The leaves have stalks. The leaves have 3 leaflets. The leaflets have short stalks. The leaflets are 2-5.5 cm long. The tendrils branch. The flowers are blue or reddish-purple. The fruit is a flat pod.

Distribution: It grows in warm temperate places. In Nepal it grows to about 1100 m altitude. In Africa it grows to 3,500 m above sea level. It grows in moist places. It is drought resistant. It grows in the highlands in Eritrea. It can grow in arid places. It suits hardiness zones 8-10.



Use: The seeds are eaten. They are used in soups. In small quantities they do not cause problems. The leaves are eaten as a pot-herb. The flour from the dried seeds is occasionally mixed with wheat or rye flour. The very young pods are cooked and eaten.

Caution: Poisonous unless treated. It can cause the legs to become paralysed. This disease called lathyrism occurs when the diet is 30-50% chickling vetch. The risk can be reduced by boiling and baking

Cultivation: Plants are grown from seed.

Production: It gives some yield even when other bean or grain crops have failed.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	8.4	1455	27.4	-	-	10.0	-

Image sourced from: www.flickr.com

Legumes

English: Rice bean

Local:

Scientific name: *Vigna umbellata*

Plant family: FABACEAE

Description: An annual twining, climbing bean plant with a slender hairy vine. It grows from seed each year. It grows to 1.5 - 3 m long. Leaves have 3 leaflets which can vary in shape. They are mostly oval and 3 - 13 cm long by 1.5 - 7 cm wide. They taper towards the tip and are rounded at the base. Usually they are hairy. The leaf stalks are 3 - 16 cm long. Flowers are about 1.5 cm long in dense cone shaped clusters. These flowering stalks can be 3 - 10 cm long. The flowers are yellow. The fruit are straight pods about 10 cm long and 5 mm wide. Seeds are small (5 - 8 mm long) and yellow to brown. The pods split open easily. The seeds can be yellow, green, brown, red, black or mottled.



Distribution: A tropical plant that grows to 1,800 m altitude in the tropics. It suits wet climates. It occasionally becomes self sown in coastal grasslands. It needs a sunny protected position and is drought and frost tender. It can grow in arid places.

Use: The young pods and ripe seeds are eaten cooked. The dried seeds are boiled and served with rice or used in soups and stews. The young leaves can be eaten. The seeds are used in bean sprouts. Seeds should be cooked or crushed if fed to pigs.

Cultivation: It is grown from seeds. Seed collection is easy. Seeds often have a hard skin which must be broken (e.g. by scraping) to help germination.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	13.0	1373	20.9	-	-	10.9	-

Leafy greens

English: Indian spinach

Local:

Scientific name: *Basella alba*

Plant family: BASELLACEAE

Description: An annual or perennial climbing herb with thick fleshy leaves. The vine is smooth and juicy and can be 10 m long. It branches freely. The vine and leaves can be red or green. The leaves are fleshy and pointed at the tip. They can be 8 - 18 cm long and 8 - 10 cm across. They are carried alternately along the vine. Leaves can be heart shaped or oval. It has white, pink or red flowers in short spikes which are in the axils of the leaves. The fruit are round and soft. They can be red, white or black and are 6 - 8 mm across. The seeds are round and black. They are 3 mm across.



(Often the ones with heart shaped leaves are called *Basella cordifolia*, the ones with a red stem *Basella rubra* and the short day flowering dark green kind *Basella alba*.)

Distribution: A tropical plant. It occurs mostly in the tropical lowlands and is best below 500 m but will grow up to about 1600 m. in the equatorial tropics. It will grow quite well in the temperature range 15 - 35°C. It does not like water-logging but can survive 4 - 12 weeks drought once well established. It requires adequate water during the growing season. The best pH is 5.5 - 7.0. It cannot tolerate salty conditions. Flowering does not occur when day lengths are over 13 hours.

Use: The young shoots and leaves are eaten cooked. They are somewhat slimy. In soups and stews the mucilage can be used as thickening. The purple colour of fruit is harmless and is used to colour vegetables and agar-agar. Some lemon juice added to the dye enhances the colour. The leaves can be eaten raw in salads or cooked like a vegetable. The leaves are used to make tea and can also be dried and stored. The seeds can be crushed to use as an edible dye for jellies.

Cultivation: It can be sown from seeds or cuttings. Seeds germinate in a few days. Sticks can be provided for support, or it can grow over fences and stumps. If seeds are used, 3 kg of seed will sow one hectare. They are best sown in a nursery and transplanted. A spacing of 1 m is suitable. Plants grown from seed are more productive than those grown from cuttings. When cuttings are used, 20 - 25 cm long cuttings are suitable. Where the plant grows over light soil it can root at the nodes and continue growing. Partial shade, rich fertile soil and adequate moisture favour abundant leaf production. It is responsive to nitrogen fertiliser. Light shade gives bigger leaves. It requires a trellis to climb over. Frequently picking of the bud encourages branching.

Production: It is 4 - 6 weeks until the first harvest. It grows reasonably well on poor soils and is fairly resistant to pest and disease. Leaves will only store for one day at 20 - 30°C. Yields of 40 kg of leaves from a 10 metre square bed is possible over 75 days.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	85.0	202	5.0	56	100	4.0	-

Leafy greens

English: Spinach

Local:

Scientific name: *Spinacia oleracea*

Plant family: AMARANTHACEAE

Description: An annual leafy vegetable. It grows 60-90 cm high and spreads 30-45 cm wide. The broad leaves are produced in a clump on short stalks. The leaves at the base are large and leaves on the stalk are smaller. Plants are separately male and female. Both types are needed if seed is to be produced. Flowers are greenish in spikes.



Distribution: It is a temperate plant. It does not suit the tropical lowlands and grows best where the temperature varies between 10°C and 20°C or above 2,000 m altitude. The kind with very prickly seeds is frost resistant. Plants need a deep well drained soil. It is a cool season, short day plant. In Nepal it grows to about 1,400 m altitude. It suits hardiness zones 6-9.

Use: Leaves are cooked in a small amount of water. They are also used in soups and salads. Young leaves are eaten raw and older leaves are cooked. The sprouted seeds can be used in salads.

Caution: Spinach can contain oxalates which affects calcium absorption.

Cultivation: It is normally sown directly from seeds. Plants need to be 25 cm apart.

Production: The older leaves are picked off. They can be harvested starting at 8 weeks.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	91.6	61	2.9	32	52	2.71	0.5
leaf (boiled)	92.9	57	2.4	819	29	2.9	0.8

Leafy greens

English: Amaranths

Local:

Scientific name: *Amaranthus tricolor*

Plant family: AMARANTHACEAE

Description: A small, annual, leafy green herb about 1 m high, spreading to 45 cm wide. An upright, much-branched annual with a thin membrane covering the stems. Sometimes the plant lies over. The stems are angular. The plant branches in the upper part of the plant. It does not have thorns and grows from seed each year. Leaves have long leaf stalks which can be 5 - 10 cm long. Leaves vary in shape, size and colour. The leaf blade can be 5 - 25 cm long by 2 - 6 cm wide. Leaves are dull-purplish and the top leaves can be yellow or red. Some types have coloured leaves or patterns on the leaves. It has a clumpy seed head at the top. The flower spike at the top can be 30 cm long. The seeds are 1 - 1.2 mm across and black.



Distribution: It grows in most tropical countries, including the Pacific and Solomon Islands. It will grow in warm, temperate places. Plants grow wild in waste places. Amaranths grow from sea level to 2,400 m altitude in the equatorial tropics. Amaranth seeds need a temperature above 15 - 17°C to germinate. In areas of the equatorial highlands above 1,800 m, average temperatures are probably below this during the cooler months. It may be more difficult to get Amaranths started during these months. It suits hardiness zones 8 - 11.

Use: The young leaves and stems are cooked and eaten as a vegetable. The seeds can be eaten. It is a very important tropical vegetable. It grows quickly, produces well and is nutritious.

Cultivation: The very small seeds of this plant are scattered over ashes or fine soil in fertile ground. The seeds are normally spread by rubbing the dry seed heads between the hands. Some types are self-sown. These plants grow in most tropical countries. The soil must be fertile. If they are put in an old garden, they will grow very poorly. They should either be put in a new garden site, when it is cleared from bush, or in old ground that is had compost added. Small gardens close to a house can often be built up to a good fertility by using food scraps and ashes that are left over near houses. Amaranths need high amounts of two nutrients, nitrogen and potash. The ashes from fires are high in potash, so farmers scatter seeds of Amaranth over areas they have burnt. Normally, the hotter it is, the better they grow. They also like plenty of sunlight and do not suit shaded places. They need to have water most of the time they are growing. This is mostly not an issue in areas with high rainfall.

Production: Plants can be harvested when small by thinning out closely-spaced plants. These can be either transplanted or eaten cooked. Plants can be harvested whole or have top leaves harvested several times. Harvesting begins after 4 - 7 weeks and can continue over the next 2 months.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	91.7	96	2.5	292	43.3	2.3	0.9

Leafy greens

English: Wild Amaranth

Local:

Scientific name: *Amaranthus blitum*

Plant family: AMARANTHACEAE

Description: An annual plant up to 1 m high. The leaf stalk is slender. The leaves are oval and 1-6 cm long. There are prominent veins under the leaf. The male and female flowers are separate but on the same plant. They are in a spike at the top of the plant. The male and female flowers are mixed. The flowers are green. The seeds are round and 1-1.8 mm across. They are dark brown to black.

Distribution: It grows in both temperate and tropical zones. It grows well with temperatures above 25°C. In Indonesia it grows from the lowlands to about 2,000 m above sea level.



Use: The leaves and young plant are eaten cooked. The seeds are ground into flour and used to make bread. An edible dye is obtained from the seed capsules.

Caution: This plant can accumulate nitrates if grown with high nitrogen inorganic fertilisers and these are poisonous.

Cultivation: Plants can be grown from seed if the soil is warm. Seeds are small and grow easily. They can be transplanted. Cuttings of growing plants root easily. The seed needs to be near the soil surface.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	-	-	3.88	92	-	8.3	-

Image sourced from: www.prota4u.org

Leafy greens

English: Hibiscus

Local:

Scientific name: *Hibiscus rosa-sinensis*

Plant family: MALVACEAE

Description: A shrubby, evergreen, woody, shrub used for hedges. It grows 2 - 5 m tall. The bark is grey and flaky and has fine stripes. The leaves are bright green and oval with long tips. The edges are entire on the lower leaves. The upper leaves are coarsely toothed. The flowers occur singly in the axils of leaves. Flowers can be single or double. They are bell shaped and 10 - 15 cm across. There are a range of colours. The fruit are rounded capsules with many seeds inside. The capsules are beaked. Plants usually do not produce fruit in the hot humid tropics.



Distribution: A tropical and subtropical plant, common as an ornamental throughout the tropics. It originally came from China. It thrives on any type of soil. Different types are adapted to sunny or shady places. It grows in open, moist places. It grows where average temperatures are 15 - 30°C. It is very sensitive to frost and can grow from sea level to 1,000 m altitude. It requires a minimum rainfall of 700 mm per year and suits hardiness zones 9 - 11.

Use: The leaves are eaten cooked. In some places they are pounded before cooking. The flowers are eaten raw or pickled. They are also added to drinks. They are used to colour foods including preserved fruit, sliced pineapple, agar-agar jellies, and cooked vegetables. The fresh flower ovary is eaten.

Cultivation: It is mostly grown from cuttings.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf (dry)	6.4	1339	25.9	-	-	19.6	8.9
leaf (fresh)	76.0	321	2.3	-	-	-	-

Leafy greens

English: Jute

Local:

Scientific name: *Corchorus olitorius*

Plant family: MALVACEAE

Description: An annual plant. It is upright, branching, and slightly woody. Plants vary in height, shape, leafiness and hairiness. Plants grown for leaves are usually only 30 cm tall. They also have many branches. Leaves are shiny and have leaf stalks. The leaves have teeth along the edge. The tips of the lowest leaves in each side, have long bristle like structures. Small clusters of yellow flowers grow in the axils of the leaves. The fruit are ridged capsules. They can be 7 cm long. These have partitions across them between the seeds. A ripe capsules contains 180 - 230 seeds. The seeds are dull grey and with four faces and one long point. Each seed has one pale line along it.



Distribution: A tropical plant. It is mostly coastal, below 250 m altitude. Temperatures of 22°-35°C are suitable. It can stand both drought (2 - 3 weeks) and water-logging, except when young. A well drained soil is best. They require humus-rich soils. A soil pH of 5.5 - 7.0 is best, but they can grow in soils with pH up to 8.5. They also need adequate moisture for good leaf production. A rainfall of 1,000 mm is suitable. A high relative humidity (80 - 90%) is best. It produces seeds when day lengths are short. It grows in most African and Asian countries.

Use: The young leaves and stem tops are eaten cooked. They are slimy unless fried. They are also used to make a thick soup. Leaves can be sun dried, pounded to flour, then stored for a long time.

Cultivation: Plants grow from seed, and they can be transplanted. Seeds are often broadcast into fine seed beds at the beginning of the wet season. Mixing the small seeds with sand makes it easier to sow them evenly. Often seeds are slow to start growing. This can be overcome by soaking them in hot water. A spacing of 20 - 30 cm between plants is suitable. For vigorous varieties this could be increases to 45 - 50 cm. Seeds are saved from pods for re-sowing.

Production: First leaves can be harvested after 5 - 6 weeks. Tips about 20 - 30 cm long are picked. Production of edible green tips, is not large. 7 - 8 kg of leaf tips can be harvested from 3 - 8 pickings over 3 - 4 months. Seeds can be collected after 13 - 15 weeks. If seeds of a particular variety are desired, it is necessary to grow these plants 16 m away from other plants, to avoid cross pollination. Seeds can be stored for 8 - 12 months in well sealed jars.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf (raw)	80.4	244	4.5	574	80	7.2	-
leaf (cooked)	87.2	155	3.4	156	33.0	3.1	0.8

Leafy greens

English: Yellow wood-sorrel

Scientific name: *Oxalis corniculata*

Local:

Plant family: OXALIDACEAE

Description: A small herb. It lies along the ground. The root system is thin. Several stems grow and these usually lie along the ground. These can form roots at the nodes. It grows 5 cm high. The stem is branched. There are no swollen storage organs under the ground. The leaves are stalked. They are alternate. The leaves have 3 leaflets. The leaflets have short slender stalks. The leaves are deeply lobed at the tip and wedge shaped at the base. There are stipules which are rounded and joined to the leaf stalk. The flowers are small and single. They occur on a side stalk. The flowers are yellow. The fruit are narrow and oblong. They are 5 angled. They have a short beak. There are several brown seeds.



Distribution: It can grow from tropical to warm temperate places. In Nepal it grows up to about 2900 m altitude. In Zimbabwe it grows between 490-2,270 m above sea level. It can grow in hot arid places. It can grow in arid places.

Use: The young leaves are chewed when fresh. They are also pickled. The leaves are sour and can be added to salads. They can be cooked as a potherb. The leaves can be soaked in hot water for 10 minutes to make a drink.

Caution: Because the plant contains oxalates, eating it over extended periods can reduce the body's ability to absorb calcium.

Cultivation: Plants are grown from seed. They can also be grown from rooted cuttings of the branches.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	85.5	188	5.1	60	98	5.2	-

Image sourced from: <http://luirig.altervista.org/photos-ni/oxalis-corniculata.htm>

Leafy greens

English: Ivy Gourd

Local:

Scientific name: *Coccinia grandis*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is a climbing or trailing herb. It can climb 20 m high. It has single tendrils. It has long tuberous roots. The leaves are heart shaped, with the lobes towards the base in an angular shape. The leaves are 5-10 cm across. The edges of the leaves are irregular. The flowers occur as male and female flowers on separate plants. The flowers are large and white and occur singly. The fruit is oblong and up to 10 cm long by 2.5 cm across. It is green with white stripes. The fruit become bright red when mature. The seeds are white or light brown. They are hairy and flattened with a broad rim.



Distribution: A tropical plant. It does well in warm climates. In cold weather, plants remain dormant. They cannot grow in waterlogged soil. Plants grow between 550 and 1600 m altitude in Ethiopia. It requires a well distributed rainfall and a fairly high humidity. It grows in areas with an annual rainfall between 1,000-1,600 mm. It requires a sandy soil with good drainage and a high level of organic material. In Nepal the plants grow up to 1400 m altitude. In South Vietnam it grows up to 1,500 m above sea level. In the Himalayas it grows between 1,200-2,000 m above sea level. It can grow in arid places.

Use: The leaves, shoots, and immature fruits are cooked as a vegetable. Mature fruits are eaten fresh. The ripe red fruit are collected, peeled and cut into small pieces and added to porridge. The leaf tips are cooked as a vegetable. The leaves can be stored for 10 days. The ripe fruit are eaten raw or cooked. They are used in curries. Sometimes the fruit are candied.

Cultivation: Plants are grown from stem cuttings or tuberous roots. They can also be grown from seeds. Stem cuttings about 12-15 cm long are suitable. The plant needs a trellis to climb over. One male plant to every ten female plants is enough to ensure pollination. A spacing of 150 cm apart is suitable. During dry periods watering is required.

Production: Plants can grow for 3 or 4 years. The yield of fruit can be about 10 kg per plant. In Tanzania the fruit are collected in June to August.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	90.6	117	4.1	36	-	4.6	-
fruit	93.6	83	0.7	-	-	0.6	-

Image sourced from: www.flickr.com

Fruit

English: Pomelo

Local:

Scientific name: *Citrus maxima*

Plant family: RUTACEAE

Description: A dome-shaped, spreading, spiny tree that grows up to 15 m tall. The glossy, oval leaves are very large. And are downy underneath. The leaf stalks have broad wings. Young shoots and stems have fine hairs on them. Flowers are large (2 cm) and creamy white. The flowers are produced in bunches from woody shoots. The flowers have a sweet scent. The fruit are oval or pear shaped. The fruit is very large (20 cm) with a thick skin. The skin is dotted with oil glands. The fruit are green but become yellow when ripe. They contain 11 - 14 segments. The flesh can be



pale yellow or pink. Each segment of the fruit is covered by a strong membrane. Some kinds have many seeds, while others are almost seedless. There are several named cultivated varieties.

Distribution: A tropical plant that thrives in warm lowland areas. It can grow from sea level up to 900 m. It is tolerant of brackish and salty conditions. It suits humid climates in tropical and subtropical locations.

Use: The fruit can be eaten fresh. It is used for desserts, jams and marmalades. The fruit pulp can be dried and candied. The skin can be eaten as a vegetable. The leaves are used to flavour meat and fish during cooking.

Cultivation: It is mostly grown from seed, and does not breed true. The seed only produce one seedling unlike many citrus. Trees are often produced by aerial layering, but budding or grafting can be used. Air-layered trees give sweeter fruit. Trees start producing after about 9 years. Trees need to be about 9 m apart. Trees grown away from other trees often produce almost seedless fruit.

Production: Fruit is produced almost all the year round. The time from flowering to ripe fruit is about 6 months. Fruit can be stored quite well.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit (raw)	90.3	175	0.6	20	37	0.4	-

Fruit

English: Pawpaw

Local:

Scientific name: *Carica papaya*

Plant family: CARICACEAE

Description: Pawpaw is a well known tropical fruit that grows 3 - 5 m tall and only occasionally has branches. The stem is softly woody and has scars from fallen leaves along it. There is a clump of leaves at the top of the plant. The leaves are large (50 cm wide) deeply lobed and on leaf stalks up to 90 cm long. Trees can be male, female or bisexual. Male flowers are small and white and on long stalks. Female and bisexual flowers are on short stalks. These have no fruit, round fruit and long fruit respectively. There are three forms of long fruit. The seeds are black.



Distribution: It is a tropical plant that grows from sea level up to about 1700 m altitude in the equatorial tropics. In cooler regions they have to be planted but in humid tropical regions are commonly self-sown. Sunlight allows germination when forest is cleared. It cannot stand frost. It needs a night temperature above 12°C and don't tolerate water-logging. Plants die after 48 hours in standing water. It needs a pH between 5 – 8 and suits hardiness zones 11 - 12.

Use: Fruit can be eaten ripe and raw. Green fruit can be cooked as a vegetable. The young leaves can be eaten cooked, but are bitter. The flowers and the middle of the stem can be eaten. Papayas contain papain which is a meat tenderiser. The dried seeds can be used as a spice.

Cultivation: Pawpaw seeds grow easily and plants grow quickly. Fresh seeds can be used. If dry seeds are used they should be soaked before planting. Seeds should be sown when temperatures are 24 - 30°C. They need a reasonably fertile soil. Seeds can be sown directly or the seeds can be put in a nursery and the seedlings transplanted. Seeds in a nursery should be sown 1 - 2 cm deep. Seedlings can be transplanted when they are about 20 cm high. Plants should be about 3 m apart. Continuous fruit production depends on fertility, temperature and moisture being adequate to maintain active growth. The fruit is produced year round but the growth and development rate decreases with temperature. The size and quality of fruit declines at lower temperatures. Pollination is by wind and insects and is not normally limiting. Normally cross and self-pollination both occur. Seeds are dispersed by birds, bats and people and remain viable for a few months.

Production: Seeds emerge in 2 - 3 weeks. Vegetative growth before flowering is 4 - 8 months. One or more fruit grow per leaf axil, about every 1 - 2 weeks under good growing conditions. With good growth, 100 fruit can be produced from one plant in a year. Pollination to maturity is about 2 - 3 months. On the coast in tropical equatorial regions, pawpaws start producing fruit after about 4 - 5 months, but in the highlands this may take 12 - 18 months. The first fruit are ready 6 - 11 months from planting. Tree life is about 2 - 3 years, although they may live for 10 - 12 years.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	75.4	378	8.0	-	140	0.77	-
fruit	88.0	163	0.5	290	54	0.4	0.18
fruit (unripe)	92.1	109	1.0	-	-	0.3	-

Fruit


English: Jackfruit

Local:

Scientific name: *Artocarpus heterophyllus*

Plant family: MORACEAE

Description: An evergreen, breadfruit family tree that grows up to 20 m tall. Trees form many branches but have one or two main trunks. The bark is smooth and dark green. Leaves of young trees have 1 or 2 lobes but mature leaves are long (15 cm) and entire. They are leathery, deep green and glossy. Flowers occur on spikes, on stalks from the trunk or main branches. Some stalks only have male flowers, others only have female flowers, while some have both male and female spikes. Generally male flowers are on short stalks among the leaves, and female flowers are on trunks.



Male flowers are 5 cm long by 2 cm wide and are dull green. Female flowers are bright green. The very large spiny fruit grow on main branches and the trunk. The fruit is a composite fruit made up from the many individual flowers of the flower cluster. Fruit can be 1 m long and weigh 36 kg. They have 6 sided fleshy spines. Each seed is surrounded by a yellow fleshy sheath. Seed are 2 - 4 cm long and 1 - 2 cm wide. There can be 100 - 500 seed in a fruit. Unlike breadfruit, there are no seedless Jackfruit. When ripe, the unopened fruit has a strong smell.

Distribution: A tropical plant that grows in the tropical lowlands and up to about 1200 m altitude. It can stand some drought, but not water-logging. Trees do best where there is year round rainfall. It yields poorly where humidity is low. It does best in a well drained, frost-free location that is warm and sunny. They are slightly more tolerant of cold than breadfruit. It suits areas with a temperature range 22 - 35°C. Trees can survive occasional frosts down to 0°C. It grows best with a soil pH of 6 - 6.5. They have some wind and salt tolerance.

Use: The flesh of ripe fruits can be eaten raw. Unripe fruit can be cooked and eaten as a vegetable. They are fried in curries, preserved in syrup, dried, cooked in milk or made into alcoholic drinks. Unripe fruit is pickled. The seeds can be boiled or roasted and eaten. Some kinds have more seeds. The young leaves and flowers are edible. They are eaten mixed with chilis, fish paste, sugar and salt.

Cultivation: Trees are usually sown from seeds, but it is normal to sow them in their final location as the plants don't transplant easily. They have a long delicate taproot which makes transplanting difficult. Fresh seed must be used (less than 4 weeks old). If fresh seeds are planted immediately they grow more quickly and more seeds germinate. It is better to use larger seed. The fleshy layer around the seeds should be removed. Seeds can be soaked in water for 24 hours to give better germination. It is best to sow seeds with the embryo pointing and the narrow end pointing downwards. Seeds germinate in 3 - 8 weeks. A spacing of 12 m is suitable. Where trees are used as a wind break, trees are spaced 6 m apart.

Air-layering can be used, and stem cuttings are also possible. Air layering is best done in the rainy season. Rooting hormones can be used to help roots develop. The shoots used for air layering should be 2 - 3 years old and brown in colour. To produce air layers, a small branch 3 - 4 cm across, is cut below a node and only part way around the stem. A ring 5 - 7 cm wide is cut and a layer of sand wrapped around the stem and covered with plastic. Using 1% IBA growth substances helps shoots to strike and form roots. Roots form in about 22 days and the stem can be cut off and planted after about 2 - 3 months. Because trees vary in their growth rate, how quickly they flower and fruit, and in the fruit quality, it is best to grow plants by using vegetative parts from good trees. This can be done by budding or grafting onto 12 month old seedlings already established in the field. Budding and grafting are not easy with jackfruit. Because the fruit develops on the trunk, early pruning to

allow 2 - 3 trunks to develop is helpful. As well, trees can be topped to prevent them becoming too tall and large.

Production: Jackfruit is a fast growing tree. It can be 7 m tall by 3 years old. Trees begin bearing after about 8 years. The fruiting season lasts about 4 months in subtropical places, but fruit can be produced year round in tropical places. Fruiting takes about 100 - 120 days from when the flower is pollinated until a fully mature fruit is developed but can take another 4 months to ripen. Trees tend to have heavy crops every second year. Pollination is by wind but hand pollination has been used to increase the amount of fruit set. Fruit can be up to 50 kg in weight. Fruit change from green to yellow when mature. Up to 250 fruit can be produced each year on large trees. Fruit does not store well but can be kept in a cool store for 4 - 6 weeks at 11 - 12°C and with high humidity.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (boiled)	57.6	673	5.0	-	10	0.7	-
fruit (raw)	74.5	395	1.5	30	6.7	0.6	0.4
leaf	75.5	360	5.0	-	-	17.5	-
seed (raw)	60.9	224	4.3	-	10	1.2	-
fruit + seed (immature)	85.8	-	1.6	-	13	0.1	-

Fruit


English: Canteloupe

Local:

Scientific name: *Cucumis melo*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is an annual climber with tendrils. It grows to 0.5 m high and spreads to 1.5 m across. The stems are soft and hairy and often angled. The leaves have lobes and often a wavy or toothed edge. They are on long leaf stalks. The leaves are often hairy underneath. The tendrils are not branched. The flowers are yellow and funnel shaped with expanded lobes. The male flowers occur in clusters and are produced before the female flowers. The fruit is round, mostly with a rough or streaky skin. It is green or yellow inside. The fruit is edible. Different kinds of melons occur. Some have a hard, warty, scaly skin. Others have a network of fine ridges over the surface.



Distribution: A tropical plant, but not suited to places with high rainfall. It suits hot dry places with a fertile well drained soil. It needs a sheltered sunny position. It is drought and frost tender. A temperature range of 24 - 28°C is best but much higher temperatures are tolerated. Mostly they are grown below 500 m altitude in the tropics. A pH of 6 - 6.7 is best. Acid soils are not suitable. It can grow in arid places. It suits hardiness zones 9 - 12.

Use: The ripe fruit are eaten raw. They are also dried, candied and made into jams, jellies and preserves. The seeds are sometimes eaten roasted. The seeds are blended with fruit juice to form a drink. Sometimes the immature fruit are cooked as a vegetable. The seeds contain an edible light oil. The young leaves are eaten as a potherb.

Cultivation: They are grown from seed planted about 1 - 4 cm deep. Plants need to be 1 - 2 m apart. Seedlings can be transplanted when about 10 - 15 cm high.

Production: Plants are ready 3 - 4 months after planting. Yields of 20 kg per 10 sq m is average.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	7.0	2319	15.8	-	-	-	-
leaf	85.0	172	4.2	72	-	-	-
fruit	93.0	109	0.5	169	30	0.4	0.2

Fruit

English: Bullock's heart

Local:

Scientific name: *Annona reticulata*

Plant family: ANNONACEAE

Description: A small tree, growing up to 7.5 m tall. It has several branches near the base. Trees lose their leaves at some times of the year. New shoots have short, brown hairs, but older wood is smooth and shiny. The leaves are long and spear-shaped, with short leaf stalks. Leaves have a clear edge. The leaves smell when crushed. Flowers are greenish-yellow. They occur in groups where the leaves join the stalk. Flowers occur on new wood growth. The fruit are reddish-brown in colour and 10 - 12 cm across. There is a fine hexagonal pattern over the fruit. Inside, there are large brown seeds. The fruit are edible, but the seeds are not eaten.



Distribution: The plant has been taken to most tropical countries. It occurs in the tropical lowlands and grows up to at least 1,200 m altitude. It can grow on poorer soils with different levels of acidity. It suits humid climates. It is less suited to dry climates. It suits hardiness zones 10 - 12.

Use: The ripe fruit is eaten fresh.

Caution: The seed kernel is poisonous.

Cultivation: Plants are normally grown from seeds. Seedling trees vary quite a bit. Seedlings are easy to transplant. A spacing of 4 - 7 m is suitable. Better kinds can be grown using budding or grafting.

Production: Trees begin fruiting at 3 - 5 years. Fruit setting is often improved by hand-pollination. Fruit can vary from 0.25 - 2.25 kg each. The fruit has 13% sugar. Fruit production is seasonal. The season is normally December to March.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	66.4	462	2.06	10	50	0.5	

Fruit

English: Bael fruit

Local:

Scientific name: *Aegle marmelos*

Plant family: RUTACEAE

Description: A medium sized tree that grows 3 - 12 m tall and spreads 2 m across. The stem is erect and thorny. The aromatic leaves are green, with 3 leaflets and generally sword shaped. It loses its leaves. The flowers are yellowish-which have a strong sweet smell. They contain both sexes and occur in clusters. The fruit is large and with a hard shell about 3 mm thick. It is 8 - 10 cm across and is yellow-green when ripe. The edible pulp is reddish or orange. The fruit is made up of small cells (about 15) each with woolly seeds.



Distribution: A tropical plant that prefers rich well drained soils in an open sunny position. It suits tropical or warm places. It appears to do best where there is a distinct dry season. It is drought and frost tender. A hot dry summer is best. It can tolerate some alkalinity and saline soils. It can grow in arid places.

Use: The fruit are eaten raw. They are also used to make drinks. The fruit are often sliced and dried. Marmalade can be made from ripe pulp. They can also be pickled or used in jams and jellies. The young shoots and leaves can be eaten raw in salad. They are also used in chutneys. The flowers are used to make a drink.

Caution: There are reports that leaves make women sterile or cause abortions.

Cultivation: It is grown from seed. Seed are taken from freshly picked ripe fruit. Seedlings are planted out after one year. It can be grown from root offshoots. They are best grown using patch budding. Trees are spaced 6 - 9 m apart. Trees can be pruned to have 4 - 6 strong branches. Suckers should be removed. It can also be grown from root suckers or air layering.

Production: It is slow growing. Trees produce in 3 - 4 years. Full production is gained after 15 years. Fruit are produced throughout the year. Fruit should be picked and not allowed to drop. There can be 200 - 400 fruit per tree. Fruit ripen in the dry season and can be ripened off the tree. Trees can continue bearing for 50 years. The fruit can be stored for 2 weeks at 30°C and for 4 months at 10°C.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	58	-	2.3	-	219	0.55	-

Fruit

English: Cape gooseberry

Local:

Scientific name: *Physalis peruviana*

Plant family: SOLANACEAE

Description: A perennial herb that grows 45 - 90 cm tall. They are often grown as annuals. It is hairy and slightly branched. The spreading branches are purplish and ribbed. The leaf blade is 6 - 15 cm long by 4 - 10 cm wide. The leaves are heart shaped at the base and taper to the tip. They are slightly wavy and toothed along the edge. The flowers occur singly and hang down in the axils of leaves. The flowers are white with violet anthers and slightly spotted petals. The fruit is a berry 1 - 1.5 cm across. They are orange-yellow or pale brown. This is inside an inflated husk. The seeds are yellow and 2 mm across. There are several named cultivated varieties.



Distribution: A temperate plant that grows in the tropical highlands. It suits warm climates and does best in warm sunny conditions. It needs well drained soil. Plants are not killed by a slight frost but it grows best free from severe frosts and strong winds. In Indonesia plants are found from 700 – 2,300 m altitude, but fruit best above 1,500 m. It can grow in arid places and suits hardiness zones 8 - 10.

Use: The ripe fruit are eaten fresh or cooked. They are used for jam. They can be dried, preserved, stewed, pureed, or used in pies, cakes, jellies and sauces. Roasted seeds are pickled. The leaves have been used instead of hops in beer. The leaves are also used as a potherb.

Cultivation: Plants are grown from seed that is broadcast over the soil. Seeds should be sown 1.5 cm deep in loose soil. Seed germinate irregularly. Plants should be spaced 45 cm apart. In the tropics, plants keep growing from year to year, but in the subtropics they regrow from seed each year. Plants can be grown from softwood cuttings from the upper parts of the shoots. Seedlings can be transplanted.

Production: Plants produce fruit in 1 year.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit (mature)	84.2	201	2.0	36	30	1.5	-

Fruit

English: Governor's plum

Local:

Scientific name: *Flacourtia indica*

Plant family: FLACOURTITACEAE

Description: A shrub or small tree that grows 5 - 15 m tall. The trunk is crooked and low branched and armed with scattered slender spines. The leaves are alternate, pointed at the base and rounded at the tip. The edges of the leaves are toothed with rounded lobes. Leaves are dark green on top and pale green underneath. They are 6 - 17 cm long and 3 - 7 cm wide. Male and female trees occur. The flowers are small and white; occur singly or in pairs in the axils of leaves or near the ends of short branches. The edible fruit are rounded, fleshy, purple or nearly black. They are smooth and about 1 cm across.



The flesh is yellowish, juicy and acid. There are 6 - 10 small flattened seeds inside.

Distribution: A tropical plant that grows in the lowlands. They thrive in dry, shrubby areas at low altitudes. Trees grow in coastal areas and up to 700 m or higher. In Africa it grows from sea level to 2,400 m above sea level. It grows in sub-tropical, broadleaved, evergreen forest. It can grow in arid places. It also grows on limestone.

Use: The fleshy pulp of the fruit is eaten raw when ripe or can be cooked and eaten or made into jelly. Fruit can be dried and stored.

Cultivation: Trees are normally grown from seed. Because the seeds have a hard seed coat it helps to scratch the seed to assist germination. Cuttings and air layering can be used. Groups of trees containing both male and female trees need to be grown from root suckers or by budding. Some kinds are self pollinating. A spacing of 12 - 16 m apart is needed.

Production: Fruit matures in 60 - 90 days from pollination.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	69.5	452	0.5	15	14	12	-

Vegetables

English: Asian Pennywort

Local:

Scientific name: *Centella asiatica*

Plant family: APIACEAE

Description: Asian pennywort is a slender perennial plant with creeping stems which root at the node. It grows to 10 cm high and spreads to 50 cm across. The stems are creeping and form roots at the nodes. Groups of leaves develop in clumps at the nodes. Leaves are round or heart shaped with a wavy edge. The edge has teeth or indentations along it. The leaves can be 8 cm across when grown in shady places. The leaves have 7 - 9 forked veins. The leaf stalks are 6 - 15 cm long. The stems and flowers are dull red or pink. The flowers are small and occur in small clusters. The fruit is round but flattened and 2 mm diameter.



Distribution: Asian pennywort is a tropical plant. It is a ground cover plant in old gardens and light rainforest. It grows best in sunny, moist, fertile places. In Papua New Guinea it grows mostly up to about 500 m altitude but will probably grow up to 2,500 m. It is drought and frost tender. It needs a temperature above 10°C. It can grow in arid places suits hardiness zones 9 - 12.

Use: The whole plant is eaten raw or cooked. The leaves are added to tossed salads or steamed and served with rice. They can be cooked in vegetable soups or stews. The plant is also used for medicine. The leaves are made into tea. The juice of the leaves diluted with water and sweetened with sugar are used as a drink.

Cultivation: New plants can be produced by seeds or by runners. Mostly plants grow and spread naturally. Cuttings or runners are the easiest way to grow the plants.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	89.3	143	1.6	-	37	3.1	-

Vegetables

English: Lotus-seed herb

Scientific name: *Alternanthera sessilis*

Local:

Plant family: AMARANTHACEAE

Description: A low lying and spreading plant which has many branches. It continues to grow from year to year. It has a strong taproot. The stem and branches are up to 60-100 cm long and near the ends there are 2 lines of hairs along the stem. The leaves are smooth and attached to the stem without a stalk. They are opposite. The leaves are 1-10 cm long and 0.2-2 cm wide. The flowers heads are white and 5-7 mm long. They grow along the plant and do not have flower stalks. It flowers all year round. The fruit are oval and compressed on the side. The seed is about 1.5 cm across. When plants are growing in water the stems become hollow and the plants float.



Distribution: A tropical plant. It grows in the lowlands and the highlands. It occurs in most tropical places. It is common in waste land at low and medium altitudes in the Philippines. It grows in open moist places from sea level to 2,000 m in Papua New Guinea. In Fiji it grows from sea level to 500 m. In Nepal it grows to 2,400 m altitude. It can grow in arid places. It is best in alkaline soil. It can grow in seasonally water logged soils and near rivers and ditches.

Use: The leaves and tender tips are cooked and eaten. They are used in soups. It is also used to prepare a cool drink. The harvested leaves can only be stored for 2-3 days.

Cultivation: It can be grown by dividing the underground stem. It can also be grown from sections which root at their nodes. It can be grown by seed.

Production: The first harvest of leaves can be taken 50-60 days after planting.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	89.3	109	4.5	57	77	-	-
shoot	-	-	5.0	577	-	16.7	-

Image sourced from: commons.wikimedia.org

Vegetables

English: Vegetable kenaf

Local:

Scientific name: *Hibiscus cannabinus*

Plant family: MALVACEAE

Description: A herb that can grow from seed each year, or keep growing from year to year. It grows up to 3.5 m tall. It has a few sharp spines. The leaf stalk is 6 - 20 cm long. The leaf blade has 2 forms. The leaves lower on the stem are heart shaped and those higher on the stem have 4 - 7 lobes arranged like fingers on a hand. These lobes are sword shaped and 2 - 12 cm long by 0.6 - 2 cm wide. They have teeth around the edge and taper at the tip. The flowers are yellow, white or ivory and red at the base. They occur singly in the axils of leaves. They are large and up to 10 cm across. They have very short stalks. The fruit is a capsule about 1.5 cm across. The seeds are kidney shaped.



Distribution: A tropical plant. It is cultivated in South China. It can grow in well-drained sandy soils and in dry but seasonally waterlogged places. It grows from 1,500 -2,100 m above sea level. It grows in areas with an annual rainfall of 500 - 635 mm. It can grow in arid places and suits hardiness zones 10 - 12. It grows in many African and Asian countries.

Use: The leaves are eaten cooked as a vegetable. They are also used as a substitute for tamarind for curries. They are used in soups. The leaves are cooked with the aid of potashes. The seeds are roasted and eaten. They are also fermented. The seeds yield an edible oil. The flowers are eaten cooked as a vegetable. The bark is sweet and is chewed by children.

Cultivation: It is usually grown from seeds but can be grown from cuttings. Seeds will last for about 8 months. Seeds germinate best at 35°C.

Production:

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	8.1	1785	20.2	-	-	-	-
leaf	79.0	280	5.5	34	-	12.1	-

Vegetables

English: Pumpkin

Local:

Scientific name: *Cucurbita maxima*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is a creeping vine with tendrils. It is an annual plant. The stems are soft and round in cross section. The leaves are large and hang loose. They are dark green and kidney shaped. The edges of the leaves are entire. There are large nodes at the base of the leaf. The tendrils are fairly stout and are divided half way along their length into many branches. Male flowers are carried on long upright stalks. The 5 petals are united into a long yellow tube. The female flowers are larger than the male and are fewer in number and carried on shorter stalks.



The fruit varies in size, colour and patterns on the skin. They can be round, oval or flattened, with yellow, orange or green skin. The surface can be smooth and rough and warty. The flesh is yellow and edible. The seeds are in the centre. The seeds are white or brown. They are flattened but plump and have a slanting scar at the top. The seeds are edible. (*C. moschata* does not have hairy stems but has fruit with a thickened stalk near where it joins the fruit.) There are a large number of cultivated varieties.

Distribution: A subtropical plant that grows from sea level to 2,400 m altitude. They need a fertile soil. *C. moschata* is better suited to coastal areas. They are frost sensitive but better suited to cooler areas than *C. moschata*. It can grow in arid places. It suits hardiness zones 8 - 11.

Use: The young leaf tips are eaten cooked. They can also be dried and stored. The fruit can be eaten cooked. They are baked, boiled, fried, steamed or mashed. They are used in pies and cakes. The seeds are edible, raw or roasted. They are also ground into a meal. The male flowers are eaten after removing the stamen and calyx.

Cultivation: They are grown from seed. Usually 2 or 3 seeds are planted together in a mound. The distance apart depends on the cultivar. Some kinds are better for leaf tips. It is good to save seed of adapted varieties.

Production: Fruit are ready for harvest after about 3 - 4 months. Seed can be saved from fruit for re-sowing but as pumpkins cross pollinate different types become mixed.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	6.9	2264	24.5	38	1.9	14.9	7.5
fruit	69.6	439	1.4	-	-	-	-
leaf	88.0	160	4.9	260	28	2.5	0.9
flower	88.7	107	1.4	173	14	0.8	0.1

Vegetables

English: Marrow

Local:

Scientific name: *Cucurbita pepo*

Plant family: CUCURBITACEAE

Description: A bristly hairy annual vine in the pumpkin family. It has branched tendrils. The stems are angular and prickly. The leaves are roughly triangular. The leaves have 5 lobes which are pointed at the end and are toothed around the edge. Male and female plants are separate on the same plant. Male flowers are carried on long grooved flower stalks. Female flowers are borne on shorter more angular stalks. The fruit stalks have furrows along them but are not fattened near the stalk. The fruit vary in shape, size and colour. Often they are oval and yellow and 20 cm long by 15 cm wide. The seeds are smaller than pumpkin and easy to separate from the tissue. The scar at their tip is rounded or horizontal, not oblique. There are a large number of cultivated varieties.



Distribution: A subtropical plant. They are more suited to drier areas. They are frost sensitive, and grow best with day temperatures between 24 - 29°C and night temperatures of 16 - 24°C. It suits tropical highland regions. It suits hardiness zones 8 - 11.

Use: The young fruit are cooked and eaten. They can be steamed, boiled or fried. They are used in pies, soups, stews and cakes. The young leaves and the ripe seeds can also be eaten cooked. The seeds are dried, salted and toasted and eaten as a snack food. The seeds can also be pressed to produce oil. The sprouted seeds are used in salads. Flowers and flower buds can be eaten boiled. They can be dried for later use.

Cultivation: They are grown from seeds. The seeds germinate after one week. They can be grown from cuttings. They are best planted on mounds. A spacing of 2 - 3 m between plants is needed. Hand pollination assists fruit setting. Plants can also be grown from cuttings as plants root at the nodes.

Production: The first usable immature fruit are ready 7 - 8 weeks after planting.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	3.7	2266	29.4	-	-	7.3	-
leaf	89.0	113	4.0	180	80	0.8	-
fruit (mature)	92.0	105	1.6	17	16	2.4	-
fruit	91.3	102	1.1	-	12	0.8	0.2
yellow fruit	92.0	97	1.0	180	8	1.4	-
immature fruit (raw)	92.0	92	1.5	-	9	0.4	0.1

Vegetables

English: Bitter cucumber

Local:

Scientific name: *Momordica charantia*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is a slender annual climber with flowers of both sexes on the one plant. It has simple tendrils and vines can be 4 m long. It has bright green lobed leaves 5 - 12 cm long on thin leaf stalks 3 - 10 cm long. The flowers have a sweet smell and 5 small, yellow petals. Fruit are green when young and orange when ripe. The fruit have a lumpy appearance, with ridges along its length and when fully ripe burst open. It has bright red covering on the seeds inside. The seeds are pale brown and 10 - 16 mm long and 7 - 10 mm wide. Considerable variation in the fruit occurs between varieties.



Distribution: A tropical plant that grows from sea level up to about 500 m and will probably grow to 1,000 m altitude in tropical regions. They require a well drained soil preferably rich in organic matter. Seeds do not germinate below 15°C. Plants grow best with temperatures of 18 - 35°C. A soil pH of 6.5 is best. It suits hardiness zones 9 - 12.

Use: The young bitter fruit are cooked and eaten. They are boiled, stuffed, fried or pickled. They are used in soups, stews and stir-fried dishes. The seed mass of the ripe fruit is used as a food flavouring. The leaves are also cooked and eaten as a flavouring. The tender shoots and leaves are sometimes eaten.

Caution: The leaves are considered to cause diarrhoea and vomiting.

Cultivation: Plants are grown from seed. For large scale plantings, 6 - 7 kg of seed are required for planting one hectare. Seeds are planted at 50 cm spacing in the place where the plants are to grow and need a stick to climb up. Often plants are grown on raised beds 2 m apart with 0.5 m between plants. The seed has a hard seed coat and germinates slowly. Soaking seeds for 24 hours before sowing gives a quicker more even germination. Regular watering is required.

Production: Fruit are ready to harvest 45 - 55 days after planting. Fruit should be harvested when young and tender. Once fruit have begun to change colour to yellow they are past maturity for eating. Early removal of young fruit also ensures continuous fruit setting. This can allow 6 - 8 successive pickings of fruit. Fruit on the plant are sometimes wrapped in paper to prevent fruit fly damage. Seed well stored can remain viable for 4 - 5 years. The young bitter fruit are cooked and eaten. The fruit is blanched or soaked in salt water to reduce the bitter taste.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	8.6	2020	18.6	-	-	-	-
leaf (raw)	84.7	252	5.0	44	170	7.1	0.3
leaf tip (boiled)	88.7	146	3.6	173	57	1.0	0.3
fruit	93.6	105	1.2	-	-	0.2	-
pod (boiled)	94.0	79	0.8	11	33	0.4	0.8
pod (raw)	94.0	71	1.0	380	84	0.4	0.8

Vegetables

English: Beetroot

Local:

Scientific name: *Beta vulgaris*

Plant family: CHENOPODIACEAE

Description: Beetroot is a dark green leafed plant up to 20 cm tall. It can be grown as an annual plant. Normally it gives a thickened root in the first year then flowers in the second year. The leaves vary in shape and colour. They can be oval with an irregular wavy edge. They can be dark green or reddish. It has a round or elongated fattened root. The root is red in colour. (White varieties also occur). The flowers are small and green and have both sexes. They occur in flower arrangements with the end bud a flower bud. This forms a tall, branching, spike-like arrangement. Often 2 or more seeds are joined together in a "seedball".



Distribution: It is a temperate climate plant. It is mainly grown in the highlands from 1,150 – 2,600 m altitude in the tropics. Plants grow best at 18 - 22°C and are frost resistant. Temperatures below 10°C cause the plant to start its seeding phase. It is sensitive to acid soils but can grow in alkaline soils up to pH 10.

Use: The red tubers are eaten after cooking. The root is also dried and powdered and the flour mixed with barley or wheat. They can be pickled or fermented as beetroot juice. They are often boiled, sliced and served with vinegar. The leaves are edible and can be cooked in soups and stews.

Cultivation: Plants are grown from seed. Normally the seeds are sown in the final site because transplanting is difficult. When the small clump of seeds or seed ball are planted more than one seedling will result. Plants get a soft heart due to boron deficiency. This is treated with borax.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
root (boiled)	82.7	189	1.8	-	5	0.4	0.4
root (raw)	87.1	118	1.3	2	6	0.8	0.4
leaf (boiled)	89	113	2.6	-	25	1.9	0.5
leaf (raw)	92	80	1.8	316	30	3.3	0.4

Nuts, seeds, herbs and other foods

English: Fat hen

Local:

Scientific name: *Chenopodium album*

Plant family: CHENOPODIACEAE

Description: An annual plant that grows to 1 m tall and spreads to 1 m across. The stem is erect and succulent with no hairs. They often have soft mealy lumps which can be rubbed off. The leaves are simple, with one at each node, and occurring alternately up the stem. The leaves are oval and wedge shaped with saw like edges. They are 5 - 12 cm long by 3 - 10 cm wide. The leaf stalk is usually shorter than the leaf blade. The under surface of the leaf often has a white mealy layer which can be rubbed off. The flowers occur in dense white spikes at the tip and ends of branches. The fruit is a small, roundish, papery pod that opens around the tip. The pod contains large numbers of shiny black seeds that are 1.2 - 1.8 mm across.



Distribution: A temperate plant that also grows in the tropics. It grows best on light to medium well drained soil. It suits an open sunny position but can tolerate shade. It is drought and frost resistant. It commonly occurs as a weed in old fields. In Zimbabwe, it grows from 1,100 - 1,600 m above sea level. It can grow in arid places and can tolerate temperatures of 5 - 30°C.

Use: The seeds can be ground into flour. They contain saponin which should be leached out. They are used for bread, pancakes, muffins and biscuits. The tender leaves are cooked and eaten as a vegetable. They are also used in stews. Young flowers are cooked and eaten. The sprouted seeds are edible.

Cultivation: Plants are grown from seed. Seedlings can be transplanted at a spacing of 30 cm. It does well in soils with lots of nitrogen. It is self sown and harvested from potato crops in India.

Production: The tops can be eaten before and after flowering. They are harvested after 40 days.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	-	1654	16	-	-	-	-
leaf (boiled)	88.9	134	3.2	391	37.0	0.7	0.3
leaf	87.7	113	5.3	33	108	-	-

Nuts, seeds, herbs and other foods

English: Lotus Root

Local:

Scientific name: *Nelumbo nucifera*

Plant family: NELUMBONACEAE

Description: A herb which grows in water and continues growing for several years. The rootstock is under water and creeping. The leaf stalk grows 1-2 m high. The leaves can be 20-90 cm across. It has large round leaves which stand out of the water. The stalk joins to the centre of the leaf. It has large attractive pink, red or white flowers which stand out of the water on long stalks. Flowers are pink or white and 15 to 25 cm across. The flower stalk can be longer than the leaf stalk. The mature fruit is a spongy cone shaped structure with several seeds about 1 cm across under the holes in the top. These fruit can be 7-15 cm long. The edible rhizome is submerge in mud.



Distribution: It is a tropical plant but will grow in cooler places. It needs 20-30°C. It needs full sunshine. It can grow in deep water. It does best in fresh water. The pH can range from 5.6-7.5. It suits hardiness zones 9-12. It is common in some parts of the Philippines such as Laguna de bay. It is also reported from Camarines, Mindoro, Cotabato and Davao. It will probably grow up to about 1000 m altitude. It occurs in the Fly and Sepik River areas in Papua New Guinea. It is cultivated in most of China except the very northern areas. It can grow in arid places. It suits hardiness zones 9-12.

Use: The seeds are eaten raw or roasted. Ripe seeds are prepared by removing the seedcoat and the intensely bitter plumule and then boiled in syrup. Slightly unripe seeds are also eaten raw. The seeds can be used as a coffee substitute. The white roots or rhizomes are eaten raw with meat. They can also be boiled, pickled, stir-fried or preserved in sugar. It is also the source of starch. The flower stalks are cooked as a vegetable. They are also eaten raw. The young tender leaves are eaten with a savoury sauce. The dry leaves are used to wrap food and impart flavour. The pollen and stamen are used for flavouring tea.

Cultivation: It can be grown from sections of the root or from seed. Seed germinate between 25-30°C. The hard seed coat needs to be cracked to enable seed to grow. Seed will remain viable for many years. Plants grown from seed vary as seed do not breed true. It needs to be in water at least 30-50 cm deep. Rhizomes with 2-3 segments are used. They are put in the mud at a 30° angle.

Production: Plants grow quickly. Rhizomes are harvested all year round and contain an abundance of starch. Plants grown from seed take 2 years to produce. Plants grown from cuttings can produce the same year.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	38.8	545	8.8	2	3	8.0	0.2
root(boiled)	81.4	276	1.6	-	27.4	0.9	0.3

Nuts, seeds, herbs and other foods

English: Peanut

Local:

Scientific name: *Arachis hypogea*

Plant family: FABACEAE

Description: Peanuts grow on spreading bushy plants up to about 40 cm high. The leaves are made up of 2 pairs of oppositely arranged leaflets. Flowers are produced in the axils of the leaves. Two main kinds of peanuts occur. The runner kind (Virginia peanut) has a vegetative or leafy branch between each fruiting branch and therefore produces a spreading bush. The bunch type (Spanish-Valencia peanuts) produces fruiting branches in a sequence one after the other along the branches. They grow as a more upright plant and grow more quickly. Pods are produced on long stalks which



extend under the ground and they contain between 2 - 6 seeds. The stalk or peg from the flower grows down into the soil and then produces the pod and seed under the ground. The flowers need to be no more than 18 cm from the soil surface for the seed pod to develop underground.

Distribution: Peanuts grow in tropical and subtropical areas. They grow well from sea level up to about 1,650 metres in the equatorial tropics. They require temperatures of 24 - 33°C. Plants are killed by frost. They need a well drained soil and cannot stand water-logging and often require raised garden beds. Peanuts need 300 - 500 mm of rain during the growing season. Dry weather is needed near harvest.

Use: The seeds can be eaten raw or cooked. They are boiled, steamed, roasted, salted or made into peanut butter or flour. The young leaves and unripe pods are edible after cooking. Sprouted seeds can be eaten. An edible oil is extracted from the seeds. The remaining meal can also be eaten.

Cultivation: Peanuts require soil with good levels of calcium and boron or they produce empty pods. Peanuts have nitrogen fixing root nodule bacteria and therefore can give good yields in soils where nitrogen is low. The nuts are normally removed from the shell before planting and are sown 2 - 3 cm deep, with 10 cm between plants and 60 - 80 cm between rows. The soil needs to be weeded and loose by the time the flowers are produced to allow the peg for the seed pods to penetrate the soil.

Production: Flowering can commence in 30 days and it takes 3.5 - 5 months until maturity. Peanuts are harvested by pulling out the plant when the top of the plants die down. After harvesting, they should be left to dry in the sun for 3 - 4 days. Virginia peanuts have a longer growing season and the seeds need to be stored for 30 days before they will start to re-grow.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	4.5	2364	24.3	-	-	2.0	3.0
seed (fresh)	45	1394	15	-	10	1.5	-
leaf	78.5	228	4.4	-	-	4.2	-

Nuts, seeds, herbs and other foods

English: Sunflower

Local:

Scientific name: *Helianthus annuus*

Plant family: ASTERACEAE

Description: An upright annual plant that varies in height from 1 - 4 m. It has a strong tap root. Plants are mostly unbranched, but may have some branches. The stems are hairy. The leaves are large and oval to heart shaped with teeth around the edges. They are roughly hairy and mid to dark green. Leaves can be 10 - 40 cm long by 5 - 20 cm wide. The leaf stalk is long. The flowers are yellow and daisy like, and 9 - 20 cm across. Sometimes they are tinged red or purple.



Distribution: A temperate plant that suits the highlands of the tropics and can stand a light frost. It needs a well drained, rich soil. It is drought and frost resistant. Sunflower grow from the equator to 55°N latitude. It does not suit the wet tropics. It cannot tolerate very acid soils. It can grow in arid places. It suits hardiness zones 4 - 11. It is widely distributed in many environments.

Use: An edible oil is extracted from the seeds and used for cooking. Sometimes seeds are eaten raw or roasted. The seeds can be ground into a meal for using in bread and cakes. They are also dried, roasted and ground and used as a coffee substitute. The seeds are boiled with water and honey to make a drink. The germinated seeds are fermented into a yogurt or cheese.

Cultivation: Plants are grown from seed. Only well filled seed should be planted. It is easy to save your own seed as dry seed stores well. A plant spacing of 1 m by 0.5 m is suitable. Seed are sown at a depth of 2 - 4 cm. Mature heads are collected by hand, dried and then threshed.

Production: Time to maturity is usually 4 - 5 months. Seeds are ready to eat when the flower starts to wither.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	5.4	2385	22.8	5	1.4	6.8	5.1

Nuts, seeds, herbs and other foods


English: Horseradish tree

Local:

Scientific name: *Moringa oleifera*

Plant family: MORINGACEAE

Description: A small, soft-wooded tree that grows 9 - 12 m tall. The tree loses its leaves during the year. The bark is grey, thick, corky and peels off in patches. The leaves are pale green and the leaf is divided 3 times. The whole leaf is 30 - 60 cm long and the leaflets are usually oval and 1 - 2 cm long. The leaflets are jointed with a gland near the joint. The flowers are pale yellow. They occur in long sprays 30 cm long. Each flower has 5 petals and of these one is erect and 4 are bent backwards. The fruit is a long capsule 30 - 100 cm long by 2 cm wide. The seed capsules are up to 45 cm long. They are roughly triangular in shape. The seeds have 3 wings. Often the fruiting kinds are grown as annual plants.



Distribution: A tropical and subtropical plant. They suit the dry lowland areas and grow up to 1,350 m altitude in the tropics. They are not hardy to frost. They cannot tolerate water-logging. A pH of 6 - 7.5 is suitable. It can grow in arid places. It suits hardiness zones 9 - 12.

Use: The young tops and leaves are eaten cooked. They are eaten as potherbs or used in soups and curries. They can be dried and stored for later use. The very young long pods are eaten cooked, especially in curries and soup. They are also pickled. The young seeds are eaten roasted or fried. Sometimes the roots are used as a horseradish substitute. A gum from the bark is used as seasoning. The bark is used for tea. The roots, leaves, flowers and fruits are eaten cooked in water and mixed with salt and chili peppers. The oil expressed from the seeds is used in salads.

Cultivation: It is best to grow plants from 1 metre long cuttings but they can be grown from seed. They can be used as a hedge and pruned regularly to produce more leaves. Properly dried seed can be stored for a long time in sealed containers in a cool place. Normally perennial types are grown from cuttings and annual types are grown from seed.

Production: Trees are fast growing. They can be pruned or topped. With one variety the tree flowers and fruits continuously while with the other variety there are flowers and fruit once per year. The fruit ripens 3 months after flowering. Annual types produce fruit 6 months after planting. Leaves are best dried in the shade to retain more of their Vitamin A.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	76.4	302	5.0	197	165	3.6	-
flower	84.2	205	3.3	-	-	5.2	-
leaf (boiled)	87	189	4.7	40	31.0	2.0	0.2
pod (raw)	88.2	155	2.1	4	141	0.4	0.5
seed	6.5	-	46.6	-	-	-	-

Nuts, seeds, herbs and other foods

English: Tamarind

Local:

Scientific name: *Tamarindus indica*

Plant family: FABACEAE

Description: A large spreading tree up to 24 m tall. It has a broad, dense, evergreen crown. The trunk can be 1 m across. The bark is rough and grey with a checkered pattern. The tree can lose its leaves in dry areas. The leaves are carried one after another along the branch. The whole leaf is 6 - 12 cm long and it is divided into 10 - 17 pairs of leaflets. These are oblong and without stalks. The whole leaf has a leaf stalk about 15 cm long. The leaflets are 1 - 2.5 cm long and 4 - 9 mm wide. They are a dull dark green with a rounded tip. The flowers are pale yellow with brown markings. The flowers are about 2.5 cm across and hang on long, many flowered stalks. The fruit is an oblong, thin-skinned, fleshy capsule. The brown seeds are inside this long rough surfaced, sausage-like fruit. This pod is 6 - 8 cm long and about 2 cm wide and contracted between the seeds. The pod cracks when mature. The seeds are shiny and hard. The edible pulp is date like and reddish brown.



Distribution: A tropical legume. The tree is cultivated in a number of coastal towns in the tropics as a street tree. It is probably best grown below 800 m altitude in the tropics. It is drought resistant and cannot stand water-logging. It does well on coastal dunes above high water level. It suits semi-arid areas. It grows in the Sahel and must be in frost free locations. In Kenya it grows from sea level to 1,600 m altitude. It suits hardiness zones 11 - 12.

Use: The pulp of the fruit is edible and is also used for drinks. The seeds are also edible when cooked. They can be roasted and ground into flour. The outer skin is removed. The young leaves, flowers and young pods are also edible and are eaten in curries. They are used to make dishes acid. They are used in sauces and chutneys. The young seedlings are also edible.

Cultivation: It can be grown by seeds or cuttings. It is best to sow seedlings in pots then transplant them, but seed can be sown direct. There are about 1,400 seeds per kg. Seed should be soaked in hot water or the seed coat nicked before sowing. Seed can be stored for 2 years if kept dry, cool and away from insects. Trees can be topped or cut back and allowed to re-grow. Nothing grows under the trees due to the acidity of the leaves. Trees can be grown by air layering or cuttings.

Production: Trees are long-lived and grow very slowly. Fruiting is seasonal from April to June. It takes 8 - 9 months from flowering to ripe fruit. If plants are grown for shoots, they are planted close together.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	38.7	995	2.3	20	60	1.1	0.7
flower	80.0	314	2.5	-	-	1.4	-
leaf	78.0	305	3.1	-	-	2.0	-

Nuts, seeds, herbs and other foods

English: Coastal almond

Local:

Scientific name: *Terminalia catappa*

Plant family: COMBRETACEAE

Description: A large tree, up to 25 - 40 m tall. It loses its leaves during the year. The trunk can be straight or twisted. There can be buttresses up to 3 m tall. The branches lie horizontally and come out in layers. The leaves are long, smooth and shiny, with an abrupt point at the tip and a rounded base. Leaves tend to be near the ends of branches. Leaves can be 17 - 29 cm long and 10 - 15 cm wide. Young leaves have soft hairs. The leaves turn red and fall off twice a year. Flowers are greenish-white and in a spike at the end of the branches. The lower flowers on a spike are female, and the others are male. The fruit is about 6 cm long by 3 - 4 cm wide, thick and flattened, with a flange around the edge. The fruit are green and turn red when ripe. The pulp is edible.



Distribution: It grows on beaches in almost all tropical countries in the world, including Solomon Islands. It is a tropical plant, and sometimes cultivated as a shade tree. The tree is common in lowland areas particularly on sandy or rocky beaches. Seeds are spread by bats and sea water, as well as being planted by people. It is common along streets in coastal towns. It will grow from sea level up to about 800 m altitude. Plants are frost-susceptible. It can tolerate drought. It suits hardiness zones 11 - 12.

Use: The kernel of the fruit is eaten raw. An edible oil can also be extracted.

Cultivation: Plants can be grown from seed. Seeds can be stored dry for a year or more. Seeds germinate freely and most seeds grow. Insects can badly damage the leaves of young seedlings.

Production: It is fast growing. Nut production is seasonal.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
nut (fresh)	31	1810	15.9	-	4	4.6	4.9
nut (dry)	4.2	2987	20.0	-	2	6.3	8.8

Nuts, seeds, herbs and other foods

English: Foxtail millet

Local:

Scientific name: *Setaria italica*

Plant family: POACEAE

Description: An annual grass. It grows 1 - 1.5 m tall. It can be tinged with purple colour. The stalks are upright and the section between the nodes is hollow. It develops tillers from the base. It has along leaf sheath. The leaf blade is 30 - 45 cm long by 1.2 - 2.5 cm wide. It has a prominent midrib and tapers towards the tip. The flower is a spike-like branching flower 7.5 - 25 cm long by 1.2 - 5 cm wide. The side branches carry 6 - 12 small spikes each with 1 - 3 bristles. The mature grain is 2 mm long. There are many named cultivated varieties.



Distribution: A warm temperate plant. It suits regions of low rainfall. It is grown from sea level to 2000 m altitude. It can tolerate a wide range of soil conditions. It cannot tolerate waterlogging or long periods of drought. It can grow in arid places.

Use: It can be cooked and eaten like rice. The seeds can be parched, popped, added to soups and sauces or made into porridge, cakes, puddings, and dumplings. The sprouted seeds can be used as a vegetable. The seeds can be used for making beer. The seeds can also be made into syrup.

Cultivation: Plants are grown by seed. Seed can be broadcast or drilled. Pure stands require 8-10 kg/ha of seed. Plants are harvested by cutting off the ears.

Production: It grows quickly. Plants mature in 80 - 120 days. Flowering occurs over 10 - 15 days. Plants can be self or cross pollinated. Yields of 800 - 900 kg/ha are common and straw yields for livestock feed can be up to 2500 kg/ha.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	13.5	1425	9.5	-	-	5.5	-

Nutritional values of food plants by plant Family

Plant Family	Scientific name	Common name	Edible part	Moisture %	Energy kJ	Protein g	Vit A µg	Vit C mg	Iron mg	Zinc mg	Page
AMARANTHACEAE	<i>Spinacia oleracea</i>	Spinach	leaf	91.6	61	2.9	32	52	2.71	0.5	32
AMARANTHACEAE	<i>Amaranthus tricolor</i>	Amaranths	leaf	91.7	96	2.5	292	43.3	2.3	0.9	33
AMARANTHACEAE	<i>Amaranthus blitum</i>	Wild Amaranth	leaf	-	-	3.88	92	-	8.3	-	34
AMARANTHACEAE	<i>Alternanthera sessilis</i>	Lotus-seed herb	leaf	89.3	109	4.5	57	77	-	-	49
ANNONACEAE	<i>Annona reticulata</i>	Bullock's heart	fruit	66.4	462	2.06	10	50	0.5	-	44
APIACEAE	<i>Centella asiatica</i>	Asian Pennywort	leaf	89.3	143	1.6	-	37	3.1	-	48
ARACEAE	<i>Colocasia esculenta</i>	Taro	root	66.8	1231	1.96	3	5	0.68	3.2	18
ASTERACEAE	<i>Guizotia abyssinica</i>	Niger seed	seed	6.2	2019	17.3	-	-	-	-	20
ASTERACEAE	<i>Helianthus annuus</i>	Sunflower	seed	5.4	2385	22.8	5	1.4	6.8	5.1	58
BASELLACEAE	<i>Basella alba</i>	Indian spinach	leaf	85.0	202	5.0	56	100	4.0	-	31
CARICACEAE	<i>Carica papaya</i>	Pawpaw	fruit	88.0	163	0.5	290	54	0.4	0.18	40
CHENOPODIACEAE	<i>Beta vulgaris</i>	Beetroot	root(boiled)	82.7	189	1.8	-	5	0.4	0.4	54
CHENOPODIACEAE	<i>Chenopodium album</i>	Fat hen	leaf(boiled)	88.9	134	3.2	391	37.0	0.7	0.3	55
COMBRETACEAE	<i>Terminalia catappa</i>	Coastal almond	nut(fresh)	31	1810	15.9	-	4	4.6	4.9	61
CONVOLVULACEAE	<i>Ipomoea batatas</i>	Sweet potato	tuber(baked)	72.9	431	1.7	961	24.6	0.5	0.3	21
CUCURBITACEAE	<i>Lagenaria siceraria</i>	Bottle gourd	fruit	93.0	88	0.5	25	10	2.4	-	23
CUCURBITACEAE	<i>Coccinia grandis</i>	Ivy Gourd	leaf	90.6	117	4.1	36	-	4.6	-	38
CUCURBITACEAE	<i>Cucumis melo</i>	Cantaloupe	fruit	93.0	109	0.5	169	30	0.4	0.2	43
CUCURBITACEAE	<i>Cucurbita maxima</i>	Pumpkin	fruit	69.6	439	1.4	-	-	-	-	51
CUCURBITACEAE	<i>Cucurbita pepo</i>	Marrow	fruit	91.3	102	1.1	-	12	0.8	0.2	52
CUCURBITACEAE	<i>Momordica charantia</i>	Bitter cucumber	fruit	93.6	105	1.2	-	-	0.2	-	53
EUPHORBIACEAE	<i>Manihot esculenta</i>	Cassava	tuber	62.8	625	1.4	30	15	0.23	0.48	16
FABACEAE	<i>Cajanus cajan</i>	Pigeon pea	seed(young boiled)	71.8	464	6.0	2	28.1	1.6	0.8	24
FABACEAE	<i>Canavalia ensiformis</i>	Jack bean	seed	10.0	1423	20.4	160	0	4.9	-	25
FABACEAE	<i>Lablab purpureus</i>	Lablab bean	seed(young)	86.9	209	3.0	14	5.1	0.8	0.4	26
FABACEAE	<i>Macrotyloma uniflora</i>	Horse gram	seed(dry)	9.7	1392	-	22.5	-	-	-	27
FABACEAE	<i>Vigna radiata</i>	Mung bean	seed	11.0	1432	22.9	55	4	7.1	-	28
FABACEAE	<i>Lathyrus sativus</i>	Chickling Vetch	seed	8.4	1455	27.4	-	-	10.0	-	19
FABACEAE	<i>Vigna umbellata</i>	Rice bean	seed	13.0	1373	20.9	-	-	10.9	-	30
FABACEAE	<i>Arachis hypogea</i>	Peanut	seed(dry)	4.5	2364	24.3	-	-	2.0	3.0	57
FABACEAE	<i>Tamarindus indica</i>	Tamarind	fruit	38.7	995	2.3	20	60	1.1	0.7	60
FLACOURTITACEAE	<i>Flacourtia indica</i>	Governor's plum	fruit	69.5	452	0.5	15	14	12	-	47
MALVACEAE	<i>Hibiscus rosa-sinensis</i>	Hibiscus	leaf(fresh)	76.0	321	2.3	-	-	-	-	35
MALVACEAE	<i>Corchorus olitorius</i>	Jute	leaf(cooked)	87.2	155	3.4	156	33.0	3.1	0.8	36

MALVACEAE	<i>Hibiscus cannabinus</i>	Vegetable kenaf	leaf	79.0	280	5.5	34	-	12.1	-	50
MORACEAE	<i>Artocarpus heterophyllus</i>	Jackfruit	fruit(raw)	74.5	395	1.5	30	6.7	0.6	0.4	41
MORINGACEAE	<i>Moringa oleifera</i>	Horseradish tree	leaf(boiled)	87	189	4.7	40	31.0	2.0	0.2	59
NELUMBONACEAE	<i>Nelumbo nucifera</i>	Lotus Root	seed	38.8	545	8.8	2	3	8.0	0.2	56
OXALIDACEAE	<i>Oxalis corniculata</i>	Yellow wood-sorrel	leaf	85.5	188	5.1	60	98	5.2	-	37
POACEAE	<i>Dactyloctenium aegyptium</i>	Comb fringe grass	seed	7.5	1234	9.8	-	-	6.9	4.7	13
POACEAE	<i>Eleusine coracana</i>	Finger millet	seed	11.7	1594	6.2	-	-	5.3	-	14
POACEAE	<i>Hordeum vulgare</i>	Pearl barley	seed(boiled)	69.6	510	2.7	-	-	6.0	-	15
POACEAE	<i>Setaria italica</i>	Foxtail millet	seed	13.5	1425	9.5	-	-	5.5	-	62
RUTACEAE	<i>Citrus maxima</i>	Pomelo	fruit (raw)	90.3	175	0.6	20	37	0.4	-	39
RUTACEAE	<i>Aegle marmelos</i>	Bael fruit	fruit	58	-	2.3	-	219	0.55	-	45
SOLANACEAE	<i>Physalis peruviana</i>	Cape gooseberry	fruit(mature)	84.2	201	2.0	36	30	1.5	-	46

*Solutions to Malnutrition
and Food Security*

**FOOD PLANT
SOLUTIONS
ROTARIAN ACTION GROUP**

www.foodplantsolutions.org